



Mellin's inverse formula

Canonical name	MellinsInverseFormula
Date of creation	2013-03-22 14:23:02
Last modified on	2013-03-22 14:23:02
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	13
Author	pahio (2872)
Entry type	Result
Classification	msc 44A10
Synonym	inverse Laplace transformation
Synonym	Bromwich integral
Synonym	Fourier-Mellin integral
Related topic	InverseLaplaceTransformOfDerivatives
Related topic	HjalmarMellin
Related topic	TelegraphEquation

It may be proven, that if a function $F(s)$ has the *inverse Laplace transform* $f(t)$, i.e. a piecewise continuous and exponentially real function f satisfying the condition

$$\mathcal{L}\{f(t)\} = F(s),$$

then $f(t)$ is uniquely determined when not regarded as different such functions which differ from each other only in a point set having Lebesgue measure zero.

The inverse Laplace transform is directly given by *Mellin's inverse formula*

$$f(t) = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{st} F(s) ds,$$

by the Finn R. H. Mellin (1854—1933). Here it must be integrated along a straight line parallel to the imaginary axis and intersecting the real axis in the point γ which must be chosen so that it is greater than the real parts of all singularities of $F(s)$.

In practice, computing the complex integral can be done by using the Cauchy residue theorem.