

planetmath.org

Math for the people, by the people.

quotients of Banach spaces by closed subspaces are Banach spaces under the quotient norm

 $Canonical\ name \qquad Quotients Of Banach Spaces By Closed Subspaces Are Banach Spaces Under The Quotients Of Banach Spaces By Closed Subspaces Are Banach Spaces Under The Quotients Of Banach Spaces By Closed Subspaces Are Banach Spaces Under The Quotients Of Banach Spaces By Closed Subspaces Are Banach Spaces Under The Quotients Of Banach Spaces By Closed Subspaces Are Banach Spaces Under The Quotients Of Banach Spaces By Closed Subspaces Are Banach Spaces Under The Quotients Of Banach Spaces Under The Under$

Date of creation 2013-03-22 17:23:01 Last modified on 2013-03-22 17:23:01 Owner asteroid (17536) Last modified by asteroid (17536)

Numerical id 7

Author asteroid (17536)

Entry type Theorem
Classification msc 46B99

Theorem - Let X be a Banach space and M a closed subspace. Then X/M with the quotient norm is a Banach space.

Proof: In to prove that X/M is a Banach space it is enough to prove that every series in X/M that converges absolutely also converges in X/M.

Let $\sum_n X_n$ be an absolutely convergent series in X/M, i.e., $\sum_n ||X_n||_{X/M} < \infty$. By definition of the quotient norm, there exists $x_n \in X_n$ such that

$$||x_n|| \le ||X_n||_{X/M} + 2^{-n}$$

It is clear that $\sum_{n} ||x_n|| < \infty$ and so, as X is a Banach space, $\sum_{n} x_n$ is convergent.

Let $x = \sum_{n} x_n$ and $s_k = \sum_{n=1}^k x_n$. We have that

$$x - s_k + M = (x + M) - (s_k + M) = (x + M) - \sum_{n=1}^k (x_n + M) = (x + M) - \sum_{n=1}^k X_n$$

Since $||x - s_k + M||_{X/M} \le ||x - s_k|| \longrightarrow 0$ we see that $\sum_n X_n$ converges in X/M to x + M. \square