



Math for the people, by the people.

locally compact groupoids

Canonical name	LocallyCompactGroupoids
Date of creation	2013-03-22 18:17:51
Last modified on	2013-03-22 18:17:51
Owner	bci1 (20947)
Last modified by	bci1 (20947)
Numerical id	45
Author	bci1 (20947)
Entry type	Topic
Classification	msc 46M20
Classification	msc 81R15
Classification	msc 46L05
Classification	msc 81R50
Classification	msc 18B40
Classification	msc 55U40
Classification	msc 22A22
Synonym	locally compact topological groupoids
Related topic	Groupoids
Related topic	TopologicalGroupoid
Related topic	LocallyCompact
Related topic	LocallyCompactHausdorffSpace
Related topic	T2Space
Related topic	SecondCountable
Related topic	QuantumGroupoids2
Related topic	GroupoidAndGroupRepresentationsRelatedToQuantumSymmetries
Related topic	GroupoidRepresentationsInducedByMeasure
Related topic	LocallyCompactHausdorffSpace
Related topic	Cat
Defines	locally compact groupoid
Defines	groupoid as a small category
Defines	topological groupoid
Defines	analytic groupoid
Defines	locally compact topological group
Defines	second countable locally compact groupoid

1 Locally compact groupoids

This is a specific topic entry defining the basics of locally compact groupoids and related concepts.

Let us first recall the related concepts of groupoid and *topological groupoid*, together with the appropriate notations needed to define a *locally compact groupoid*.

1.0.1 Groupoids and topological groupoids: categorical definitions

Recall that a groupoid \mathbf{G} is a small category with inverses over its set of objects $X = \text{Ob}(\mathbf{G})$. One writes \mathbf{G}_x^y for the set of morphisms in \mathbf{G} from x to y .

A *topological groupoid* consists of a space \mathbf{G} , a distinguished subspace $\mathbf{G}^{(0)} = \text{Ob}(\mathbf{G}) \subset \mathbf{G}$, called *the space of objects* of \mathbf{G} , together with maps

$$r, s : \mathbf{G} \rightrightarrows \mathbf{G}^{(0)} \quad (1.1)$$

called the *range* and *source maps* respectively, together with a law of composition

$$\circ : \mathbf{G}^{(2)} := \mathbf{G} \times_{\mathbf{G}^{(0)}} \mathbf{G} = \{ (\gamma_1, \gamma_2) \in \mathbf{G} \times \mathbf{G} : s(\gamma_1) = r(\gamma_2) \} \longrightarrow \mathbf{G}, \quad (1.2)$$

such that the following hold :

- (1) $s(\gamma_1 \circ \gamma_2) = r(\gamma_2)$, $r(\gamma_1 \circ \gamma_2) = r(\gamma_1)$, for all $(\gamma_1, \gamma_2) \in \mathbf{G}^{(2)}$.
- (2) $s(x) = r(x) = x$, for all $x \in \mathbf{G}^{(0)}$.
- (3) $\gamma \circ s(\gamma) = \gamma$, $r(\gamma) \circ \gamma = \gamma$, for all $\gamma \in \mathbf{G}$.
- (4) $(\gamma_1 \circ \gamma_2) \circ \gamma_3 = \gamma_1 \circ (\gamma_2 \circ \gamma_3)$.
- (5) Each γ has a two-sided inverse γ^{-1} with $\gamma\gamma^{-1} = r(\gamma)$, $\gamma^{-1}\gamma = s(\gamma)$.

Furthermore, only for topological groupoids the inverse map needs be continuous. It is usual to call $\mathbf{G}^{(0)} = \text{Ob}(\mathbf{G})$ *the set of objects* of \mathbf{G} . For $u \in \text{Ob}(\mathbf{G})$, the set of arrows $u \longrightarrow u$ forms a group \mathbf{G}_u , called the *isotropy group* of \mathbf{G} at u .

Thus, as is well known, a topological groupoid is just a groupoid internal to the <http://planetmath.org/Cat> of topological spaces and continuous maps. The notion of internal groupoid has proved significant in a number of fields, since groupoids generalize bundles of groups, group actions, and equivalence relations. For a further study of groupoids we refer the reader to ref. [?].

1.1 Locally compact and analytic groupoids

Definition 1.1. A *locally compact groupoid* \mathbf{G}_{lc} is defined as a groupoid that has also the topological structure of a second countable, <http://planetmath.org/LocallyCompactHausdorffSpace> locally compact Hausdorff space, and if the product and also inversion maps are continuous. Moreover, each \mathbf{G}_{lc}^u as well as the unit space \mathbf{G}_{lc}^0 is closed in \mathbf{G}_{lc} .

Remark 1.1. The locally compact Hausdorff second countable spaces are *analytic*.

One can therefore say also that \mathbf{G}_{lc} is analytic.

When the groupoid \mathbf{G}_{lc} has only one object in its object space, that is, when it becomes a group, the above definition is restricted to that of a *locally compact topological group*; it is then a special case of a one-object category with all of its morphisms being invertible, that is also endowed with a locally compact, topological structure.

References

- [1] R. Brown. (2006). *Topology and Groupoids*. BookSurgeLLC