



Math for the people, by the people.

compact operator

Canonical name	CompactOperator
Date of creation	2013-03-22 14:26:59
Last modified on	2013-03-22 14:26:59
Owner	mhale (572)
Last modified by	mhale (572)
Numerical id	8
Author	mhale (572)
Entry type	Definition
Classification	msc 46B99
Synonym	completely continuous

Let X and Y be two Banach spaces. A **compact operator** (completely continuous operator) is a linear operator $T: X \rightarrow Y$ that maps the unit ball in X to a set in Y with compact closure. It can be shown that a compact operator is necessarily a bounded operator.

The set of all compact operators on X , commonly denoted by $\mathbb{K}(X)$, is a closed two-sided ideal of the set of all bounded operators on X , $\mathbb{B}(X)$.

Any bounded operator which is the norm limit of a sequence of finite rank operators is compact. In the case of Hilbert spaces, the converse is also true. That is, any compact operator on a Hilbert space is a norm limit of finite rank operators.

Example 1 (Integral operators)

Let $k(x, y)$, with $x, y \in [0, 1]$, be a continuous function. The operator defined by

$$(T\psi)(x) = \int_0^1 k(x, y)\psi(y) \, dy, \quad \psi \in C([0, 1])$$

is compact.