



planetmath.org

Math for the people, by the people.

Banach \*-algebra representation

Canonical name	BanachalgebraRepresentation
Date of creation	2013-03-22 17:27:37
Last modified on	2013-03-22 17:27:37
Owner	asteroid (17536)
Last modified by	asteroid (17536)
Numerical id	23
Author	asteroid (17536)
Entry type	Definition
Classification	msc 46H15
Classification	msc 46K10
Defines	subrepresentation
Defines	cyclic representation
Defines	cyclic vector
Defines	nondegenerate representation
Defines	topologically irreducible
Defines	algebraically irreducible
Defines	direct sum of representations
Defines	unitarily equivalent

## Definition:

A **representation** of a Banach  $*$ -algebra  $\mathcal{A}$  is a  $*$ -homomorphism  $\pi : \mathcal{A} \longrightarrow \mathcal{B}(H)$  of  $\mathcal{A}$  into the  $*$ -algebra of bounded operators on some Hilbert space  $H$ .

The set of all representations of  $\mathcal{A}$  on a Hilbert space  $H$  is denoted  $rep(\mathcal{A}, H)$ .

## Special kinds of representations:

- A **subrepresentation** of a representation  $\pi \in rep(\mathcal{A}, H)$  is a representation  $\pi_0 \in rep(\mathcal{A}, H_0)$  obtained from  $\pi$  by restricting to a closed  $\pi(\mathcal{A})$ -invariant subspace<sup>1</sup>  $H_0 \subseteq H$ .
- A representation  $\pi \in rep(\mathcal{A}, H)$  is said to be **nondegenerate** if one of the following equivalent conditions hold:
  1.  $\pi(x)\xi = 0 \quad \forall x \in \mathcal{A} \implies \xi = 0$ , where  $\xi \in H$ .
  2.  $H$  is the closed linear span of the set of vectors  $\pi(\mathcal{A})H := \{\pi(x)\xi : x \in \mathcal{A}, \xi \in H\}$
- A representation  $\pi \in rep(\mathcal{A}, H)$  is said to be **topologically irreducible** (or just ) if the only closed  $\pi(\mathcal{A})$ -invariant of  $H$  are the trivial ones,  $\{0\}$  and  $H$ .
- A representation  $\pi \in rep(\mathcal{A}, H)$  is said to be **algebraically irreducible** if the only  $\pi(\mathcal{A})$ -invariant of  $H$  (not necessarily closed) are the trivial ones,  $\{0\}$  and  $H$ .
- Given two representations  $\pi_1 \in rep(\mathcal{A}, H_1)$  and  $\pi_2 \in rep(\mathcal{A}, H_2)$ , the of  $\pi_1$  and  $\pi_2$  is the representation  $\pi_1 \oplus \pi_2 \in rep(\mathcal{A}, H_1 \oplus H_2)$  given by  $\pi_1 \oplus \pi_2(x) := \pi_1(x) \oplus \pi_2(x)$ ,  $x \in \mathcal{A}$ .

More generally, given a family  $\{\pi_i\}_{i \in I}$  of representations, with  $\pi_i \in rep(\mathcal{A}, H_i)$ , their is the representation  $\bigoplus_{i \in I} \pi_i \in rep(\mathcal{A}, \bigoplus_{i \in I} H_i)$ , in the direct sum of Hilbert spaces  $\bigoplus_{i \in I} H_i$ , such that  $(\bigoplus_{i \in I} \pi_i)(x) := \bigoplus_{i \in I} \pi_i(x)$  is the <http://planetmath.org/DirectSumOfBoundedOperatorsOnHilbertSpace> sum of the family of bounded operators  $\{\pi_i(x)\}_{i \in I}$ .

---

<sup>1</sup>by a  $\pi(\mathcal{A})$ - we a subspace which is invariant under every operator  $\pi(a)$  with  $a \in \mathcal{A}$

- Two representations  $\pi_1 \in \text{rep}(\mathcal{A}, H_1)$  and  $\pi_2 \in \text{rep}(\mathcal{A}, H_2)$  of a Banach \*-algebra  $\mathcal{A}$  are said to be **unitarily equivalent** if there is a unitary  $U : H_1 \longrightarrow H_2$  such that

$$\pi_2(a) = U\pi_1(a)U^* \quad \forall a \in \mathcal{A}$$

- A representation  $\pi \in \text{rep}(\mathcal{A}, H)$  is said to be **cyclic** if there exists a vector  $\xi \in H$  such that the set

$$\pi(\mathcal{A})\xi := \{\pi(a)\xi : a \in \mathcal{A}\}$$

is dense in  $H$ . Such a vector is called a **cyclic vector** for the representation  $\pi$ .

Linked file: <http://aux.planetmath.org/files/objects/9843/BanachAlgebraRepresentation.pdf>