

planetmath.org

Math for the people, by the people.

 C^n

Canonical name Cn

Date of creation 2013-03-22 14:59:43 Last modified on 2013-03-22 14:59:43

Owner drini (3) Last modified by drini (3)

Numerical id 13

Author drini (3)
Entry type Definition
Classification msc 46G05
Classification msc 26B05
Classification msc 26A99
Classification msc 26A24
Classification msc 26A15

 $\begin{array}{ccc} \text{Synonym} & & C^1 \\ \text{Synonym} & & C^2 \\ \text{Synonym} & & C^k \\ \text{Synonym} & & C^{\infty} \end{array}$

Related topic Derivative

Related topic SmoothFunctionsWithCompactSupport

Let $f: \mathbb{R} \to \mathbb{R}$ be a function. We say that f is of class C^1 if f' exists and is continuous.

We also say that f is of class C^n if its n-th derivative exists and is continuous (and therefore all other previous derivatives exist and are continuous too).

The class of continuous functions is denoted by C^0 . So we get the following relationship among these classes:

$$C^0 \supset C^1 \supset C^2 \supset C^3 \supset \dots$$

Finally, the class of functions that have continuous derivatives of any order is denoted by C^{∞} and thus

$$C^{\infty} = \bigcap_{n=0}^{\infty} C^n.$$

entry). Therefore, $f \in C^{\infty}$ if and only if every derivative of f exists. The previous concepts can be extended to functions $f: \mathbb{R}^m \to \mathbb{R}$, where f

It holds that any function that is differentiable is also continuous (see http://planetmath.org/Dif

The previous concepts can be extended to functions $f: \mathbb{R}^m \to \mathbb{R}$, where f being of class C^n amounts to asking that all the partial derivatives of order n be continuous. For instance, $f: \mathbb{R}^m \to \mathbb{R}$ being C^2 means that

$$\frac{\partial^2 f}{\partial x_i \partial x_i}$$

exists and are all continuous for any i, j from 1 to m.

C^n functions on an open set of \mathbb{R}^m

Sometimes we need to talk about continuity not globally on \mathbb{R} , but on some interval or open set.

If $U \subseteq \mathbb{R}^m$ is an open set, and $f: U \to \mathbb{R}$ (or $f: U \to \mathbb{C}$) we say that f is of class C^n if $\partial^{\alpha} f$ exist and are continuous for all multi-indices α with $|\alpha| \leq n$. See http://planetmath.org/MultiIndexNotationthis page for the multi-index notation.