



planetmath.org

Math for the people, by the people.

properties of the transpose operator

Canonical name	PropertiesOfTheTransposeOperator
Date of creation	2013-03-22 17:36:02
Last modified on	2013-03-22 17:36:02
Owner	asteroid (17536)
Last modified by	asteroid (17536)
Numerical id	7
Author	asteroid (17536)
Entry type	Theorem
Classification	msc 46-00
Classification	msc 47A05
Synonym	transpose operator properties

In this entry, whenever V, W are normed vector spaces, $\mathcal{B}(V, W)$ denotes the algebra of bounded linear operators $V \longrightarrow W$.

Let X, Y, Z be normed vector spaces and X', Y', Z' be their continuous dual spaces. Let $T, S \in \mathcal{B}(X, Y)$, $R \in \mathcal{B}(Y, Z)$ and $\lambda \in \mathbb{C}$.

0.0.1 Basic properties

- $T' \in \mathcal{B}(Y', X')$ and $\|T\| = \|T'\|$.
- $(\lambda T)' = \lambda T'$.
- $(S + T)' = S' + T'$.
- $(RT)' = T'R'$.
- If T^{-1} exists and $T^{-1} \in \mathcal{B}(Y, X)$ then $(T')^{-1} \in \mathcal{B}(X', Y')$ and $(T')^{-1} = (T^{-1})'$.

0.0.2 Miscellaneous properties

- If we endow X' and Y' with the weak-* topology then $T' : Y' \longrightarrow X'$ is continuous.
- T is an isometric isomorphism if and only if T' is an isometric isomorphism.
- If T is <http://planetmath.org/CompactOperatorcompact> then T' is also compact.
- If T' is compact and Y is a Banach space, then T is also compact.