



Math for the people, by the people.

commutant

Canonical name	Commutant
Date of creation	2013-03-22 17:21:53
Last modified on	2013-03-22 17:21:53
Owner	asteroid (17536)
Last modified by	asteroid (17536)
Numerical id	11
Author	asteroid (17536)
Entry type	Definition
Classification	msc 46L10
Related topic	VonNeumannAlgebra
Defines	double commutant

Definition

Let H be an Hilbert Space, $B(H)$ the algebra of bounded operators in H and $\mathcal{F} \subset B(H)$.

The **commutant** of \mathcal{F} , usually denoted \mathcal{F}' , is the subset of $B(H)$ consisting of all elements that commute with every element of \mathcal{F} , that is

$$\mathcal{F}' = \{T \in B(H) : TS = ST, \quad \forall S \in \mathcal{F}\}$$

The **double commutant** of \mathcal{F} is just $(\mathcal{F}')'$ and is usually denoted \mathcal{F}'' .

Properties:

- If $\mathcal{F}_1 \subseteq \mathcal{F}_2$, then $\mathcal{F}_2' \subseteq \mathcal{F}_1'$.
- $\mathcal{F} \subseteq \mathcal{F}''$.
- If \mathcal{A} is a subalgebra of $B(H)$, then $\mathcal{A} \cap \mathcal{A}'$ is the <http://planetmath.org/CenterOfARingcentralizer> of \mathcal{A} .
- If \mathcal{F} is self-adjoint then \mathcal{F}' is self-adjoint.
- \mathcal{F}' is always a subalgebra of $B(H)$ that contains the identity operator and is closed in the weak operator topology.
- If \mathcal{F} is self-adjoint then \mathcal{F}' is a von Neumann algebra.

Remark: The commutant is a particular case of the more general definition of centralizer.