



Math for the people, by the people.

## Sobolev space

Canonical name	SobolevSpace
Date of creation	2013-03-22 14:54:55
Last modified on	2013-03-22 14:54:55
Owner	paolini (1187)
Last modified by	paolini (1187)
Numerical id	10
Author	paolini (1187)
Entry type	Definition
Classification	msc 46E35
Synonym	Sobolev function
Related topic	WeakDerivative

We define the Sobolev spaces of functions  $W^{m,p}(\Omega)$  where  $\Omega$  is an open subset of  $\mathbf{R}^n$ ,  $m \geq 0$  is an integer and  $p \in [1, +\infty]$ .

The spaces  $W^{0,p}(\Omega)$  are simply defined to be the spaces  $L^p(\Omega)$  of Lebesgue  $p$ -summable functions. We then define the space  $W^{m,p}(\Omega)$  to be the space of functions  $u \in L^p(\Omega)$  which have weak derivatives  $g = (g_1, \dots, g_n)$  such that  $g_i \in W^{m-1,p}(\Omega)$ .

The space  $W^{m,p}$  turns out to be a Banach space when endowed with the norm

$$\|u\|_{W^{m,p}} = \sum_{k=0}^m \sum_{i_1=1}^n \cdots \sum_{i_k=1}^n \left[ \int_{\Omega} \left| \frac{\partial^k u(x)}{\partial x_{i_1} \cdots \partial x_{i_k}} \right|^p dx \right]^{\frac{1}{p}}$$

i.e. the sum of the  $L^p$  norms of  $u$  and of all weak derivatives of  $u$  up to the  $m$ -th order.

Of particular interest are the spaces  $H^m(\Omega) := W^{m,2}(\Omega)$  which turn out to be Hilbert spaces with the scalar product given by

$$(u, v)_{H^m(\Omega)} = \sum_{k=0}^m \sum_{i_1=1}^n \cdots \sum_{i_k=1}^n \int_{\Omega} \frac{\partial^k u(x)}{\partial x_{i_1} \cdots \partial x_{i_k}} \frac{\partial^k v(x)}{\partial x_{i_1} \cdots \partial x_{i_k}} dx.$$