

planetmath.org

Math for the people, by the people.

von Neumann algebra

Canonical name VonNeumannAlgebra
Date of creation 2013-03-22 17:21:44
Last modified on Owner asteroid (17536)
Last modified by asteroid (17536)

Numerical id 29

Author asteroid (17536)

Entry type Definition
Classification msc 46C15Classification msc 46H35Classification msc 46L10Synonym W^* -algebra
Related topic CAlgebra

Related topic TopologicalAlgebra

Related topic Commutant

Related topic GroupoidCDynamicalSystem

Related topic Algebras2
Related topic CAlgebra3

Related topic WeakHopfCAlgebra2

Related topic HAlgebra

Related topic LocallyCompactQuantumGroup

Related topic QuantumGroupsAndVonNeumannAlgebras

Definition

Let H be an Hilbert space, and let B(H) be the *-algebra of bounded operators in H.

A von Neumann algebra (or W^* -algebra) \mathcal{M} is a *-subalgebra of B(H) that contains the identity operator and satisfies one of the following equivalent conditions:

- 1. \mathcal{M} is closed in the weak operator topology.
- 2. \mathcal{M} is closed in the strong operator topology.
- 3. $\mathcal{M} = \mathcal{M}''$, i.e. \mathcal{M} equals its double commutant.

The equivalence between the above conditions is given by the von Neumann double commutant theorem.

Since the weak and strong operator topology are weaker than the norm topology, it follows that every von Neumann algebra is a norm closed *-subalgebra of B(H). Thus, von Neumann algebras are a particular class of http://planetmath.org/CAlgebra C^* -algebras and the results and tools from the C^* theory are also applicable in the setting of von Neumann algebras. Nevertheless, the philosophy behind von Neumann algebras is quite different from that of C^* -algebras and the tools and techniques for each theory turn out to be different as well.

Examples:

- 1. B(H) is itself a von Neumann algebra.
- 2. http://planetmath.org/LinftyXDmu $L^{\infty}(\mathbb{R})$ as subalgebra of $B(L^2(\mathbb{R}))$ is a von Neumann algebra.