



Math for the people, by the people.

## Fredholm operator

Canonical name	FredholmOperator
Date of creation	2013-03-22 12:58:52
Last modified on	2013-03-22 12:58:52
Owner	mhale (572)
Last modified by	mhale (572)
Numerical id	15
Author	mhale (572)
Entry type	Definition
Classification	msc 47A53
Related topic	FredholmIndex

A **Fredholm operator** is a bounded operator between Banach spaces that has a finite dimensional kernel and cokernel (and closed range). Equivalently, it is invertible modulo compact operators. That is, if  $F: X \rightarrow Y$  is a Fredholm operator between two vector spaces  $X$  and  $Y$ , then there exists a bounded operator  $G: Y \rightarrow X$  such that

$$GF - \mathbb{I}_X \in \mathbb{K}(X), \quad FG - \mathbb{I}_Y \in \mathbb{K}(Y), \quad (1)$$

where  $\mathbb{K}(X)$  denotes the space of compact operators on  $X$ . (Another way to say this is that  $F$  is invertible in the Calkin algebra). The set of Fredholm operators  $\{F: X \rightarrow X\}$  is an open subset of the Banach algebra of bounded operators  $\{T: X \rightarrow X\}$ .

If  $F$  is Fredholm then so is its adjoint,  $F^*$ . If  $T \in \mathbb{K}(X, Y)$  is a compact operator then  $F + T$  is also Fredholm.