

planetmath.org

Math for the people, by the people.

operator norm

Canonical name OperatorNorm

Date of creation 2013-03-22 12:43:20

Last modified on 2013-03-22 12:43:20

Owner asteroid (17536)

Last modified by asteroid (17536)

Numerical id 15

Author asteroid (17536)

Entry type Definition
Classification msc 47L25
Classification msc 46A32
Classification msc 47A30
Synonym induced norm
Related topic VectorNorm

Related topic OperatorTopologies

Related topic HomomorphismsOfCAlgebrasAreContinuous

Related topic CAlgebra

Defines bounded linear map
Defines unbounded linear map
Defines bounded operator
Defines unbounded operator

Definition

Let $A: V \to W$ be a linear map between normed vector spaces V and W. To each such map (operator) A we can assign a non-negative number $||A||_{\text{op}}$ defined by

$$||A||_{\mathrm{op}} := \sup_{\substack{\mathbf{v} \in V \\ \mathbf{v} \neq \mathbf{0}}} \frac{||A\mathbf{v}||}{||\mathbf{v}||},$$

where the supremum $||A||_{op}$ could be finite or infinite. Equivalently, the above definition can be written as

$$||A||_{\mathrm{op}} := \sup_{\mathbf{v} \in \mathsf{V}} ||A\mathbf{v}|| = \sup_{\mathbf{v} \in \mathsf{V}} ||A\mathbf{v}||.$$
$$||\mathbf{v}|| = 1 \qquad 0 < ||\mathbf{v}|| < 1$$

By convention, if V is the zero vector space, any operator from V to W must be the zero operator and is assigned zero norm.

 $||A||_{\text{op}}$ is called the the *operator norm* (or the *induced norm*) of A, for reasons that will be clear in the next.

Operator norm is in fact a norm

Definition - If $||A||_{\text{op}}$ is finite, we say that A is a . Otherwise, we say that A is .

It turns out that, for bounded operators, $\|\cdot\|_{op}$ satisfies all the properties of a norm (hence the name *operator norm*). The proof follows immediately from the definition:

Positivity: Since $||A\mathbf{v}|| \ge 0$, by definition $||A||_{\text{op}} \ge 0$. Also, $||A\mathbf{v}|| = 0$ identically only if A = 0. Hence $||A||_{\text{op}} = 0$ only if A = 0.

Absolute homogeneity: Since $\|\lambda A \mathbf{v}\| = |\lambda| \|A \mathbf{v}\|$, by definition $\|\lambda A\|_{\text{op}} = \|\lambda| \|A\|_{\text{op}}$.

Triangle inequality: Since $||(A+B)\mathbf{v}|| = ||A\mathbf{v} + B\mathbf{v}|| \le ||A\mathbf{v}|| + ||B\mathbf{v}||$, by definition $||A+B||_{\text{op}} \le ||A||_{\text{op}} + ||B||_{\text{op}}$.

The set L(V, W) of bounded linear maps from V to W forms a vector space and $\|\cdot\|_{op}$ defines a norm in it.

Example

Suppose that $\mathsf{V} = (\mathbb{R}^n, \|\cdot\|_p)$ and $\mathsf{W} = (\mathbb{R}^n, \|\cdot\|_p)$, where $\|\cdot\|_p$ is the vector p-norm. Then the operator norm $\|\cdot\|_{\mathrm{op}} = \|\cdot\|_p$ is the matrix p-norm.