



planetmath.org

Math for the people, by the people.

proof of Brouwer fixed point theorem

|                  |                                 |
|------------------|---------------------------------|
| Canonical name   | ProofOfBrouwerFixedPointTheorem |
| Date of creation | 2013-03-22 13:11:24             |
| Last modified on | 2013-03-22 13:11:24             |
| Owner            | bwebste (988)                   |
| Last modified by | bwebste (988)                   |
| Numerical id     | 6                               |
| Author           | bwebste (988)                   |
| Entry type       | Proof                           |
| Classification   | msc 47H10                       |
| Classification   | msc 54H25                       |
| Classification   | msc 55M20                       |

Proof of the Brouwer fixed point theorem:

Assume that there does exist a map from  $f : B^n \rightarrow B^n$  with no fixed point. Then let  $g(x)$  be the following map: Start at  $f(x)$ , draw the ray going through  $x$  and then let  $g(x)$  be the first intersection of that line with the sphere. This map is continuous and well defined only because  $f$  fixes no point. Also, it is not hard to see that it must be the identity on the boundary sphere. Thus we have a map  $g : B^n \rightarrow S^{n-1}$ , which is the identity on  $S^{n-1} = \partial B^n$ , that is, a retraction. Now, if  $i : S^{n-1} \rightarrow B^n$  is the inclusion map,  $g \circ i = \text{id}_{S^{n-1}}$ . Applying the reduced homology functor, we find that  $g_* \circ i_* = \text{id}_{\tilde{H}_{n-1}(S^{n-1})}$ , where  $*$  indicates the induced map on homology.

But, it is a well-known fact that  $\tilde{H}_{n-1}(B^n) = 0$  (since  $B^n$  is contractible), and that  $\tilde{H}_{n-1}(S^{n-1}) = \mathbb{Z}$ . Thus we have an isomorphism of a non-zero group onto itself factoring through a trivial group, which is clearly impossible. Thus we have a contradiction, and no such map  $f$  exists.