



planetmath.org

Math for the people, by the people.

adjoint

Canonical name	Adjoint
Date of creation	2013-03-22 13:48:09
Last modified on	2013-03-22 13:48:09
Owner	Koro (127)
Last modified by	Koro (127)
Numerical id	10
Author	Koro (127)
Entry type	Definition
Classification	msc 47A05
Synonym	adjoint operator
Related topic	TransposeOperator

Let \mathcal{H} be a Hilbert space and let $A: \mathcal{D}(A) \subset \mathcal{H} \rightarrow \mathcal{H}$ be a densely defined linear operator. Suppose that for some $y \in \mathcal{H}$, there exists $z \in \mathcal{H}$ such that $(Ax, y) = (x, z)$ for all $x \in \mathcal{D}(A)$. Then such z is unique, for if z' is another element of \mathcal{H} satisfying that condition, we have $(x, z - z') = 0$ for all $x \in \mathcal{D}(A)$, which implies $z - z' = 0$ since $\mathcal{D}(A)$ is dense. Hence we may define a new operator $A^*: \mathcal{D}(A^*) \subset \mathcal{H} \rightarrow \mathcal{H}$ by

$$\begin{aligned}\mathcal{D}(A^*) &= \{y \in \mathcal{H} : \text{there is } z \in \mathcal{H} \text{ such that } (Ax, y) = (x, z)\}, \\ A^*(y) &= z.\end{aligned}$$

It is easy to see that A^* is linear, and it is called the **adjoint** of A .

Remark. The requirement for A to be densely defined is essential, for otherwise we cannot guarantee A^* to be well defined.