



Math for the people, by the people.

## invariant subspace problem

Canonical name	InvariantSubspaceProblem
Date of creation	2013-03-22 17:24:02
Last modified on	2013-03-22 17:24:02
Owner	asteroid (17536)
Last modified by	asteroid (17536)
Numerical id	5
Author	asteroid (17536)
Entry type	Conjecture
Classification	msc 47A15
Classification	msc 46-00
Synonym	invariant subspace conjecture

Initially formulated for Banach spaces, the **invariant subspace conjecture** stated the following:

*Let  $X$  be a complex Banach space. Then every bounded operator  $T$  in  $X$  has a non-trivial <http://planetmath.org/ClosedSet>closed invariant subspace, i.e. there exists a closed vector subspace  $S \subset X$  such that  $S \neq 0$ ,  $S \neq X$  and  $T(S) \subseteq S$ .*

This conjecture was proven to be false when P. Enflo (1975) and . Read (1984) gave examples of bounded operators which did not have the above property.

However, if one considers only Hilbert spaces, this is still an open problem. Today the **invariant subspace conjecture** is formulated as follows:

*Let  $H$  be a complex Hilbert space. Then every bounded operator  $T$  in  $H$  has a non-trivial invariant subspace, i.e. there exists a closed vector subspace  $S \subset H$  such that  $S \neq 0$ ,  $S \neq H$  and  $T(S) \subseteq S$ .*