



Math for the people, by the people.

bounded inverse theorem

Canonical name	BoundedInverseTheorem
Date of creation	2013-03-22 17:30:51
Last modified on	2013-03-22 17:30:51
Owner	asteroid (17536)
Last modified by	asteroid (17536)
Numerical id	11
Author	asteroid (17536)
Entry type	Corollary
Classification	msc 47A05
Classification	msc 46A30
Synonym	inverse mapping theorem

The next result is a corollary of the open mapping theorem. It is often called the **bounded inverse theorem** or the **inverse mapping theorem**.

**Theorem -** Let  $X, Y$  be Banach spaces. Let  $T : X \longrightarrow Y$  be an invertible bounded operator. Then  $T^{-1}$  is also .

**Proof :**  $T$  is a surjective continuous operator between the Banach spaces  $X$  and  $Y$ . Therefore, by the open mapping theorem,  $T$  takes open sets to open sets.

So, for every open set  $U \subseteq X$ ,  $T(U)$  is open in  $Y$ .

Hence  $(T^{-1})^{-1}(U)$  is open in  $Y$ , which proves that  $T^{-1}$  is continuous, i.e. bounded.  $\square$

### 0.0.1 Remark

It is usually of great importance to know if a bounded operator  $T : X \longrightarrow Y$  has a bounded inverse. For example, suppose the equation

$$Tx = y$$

has unique solutions  $x$  for every given  $y \in Y$ . Suppose also that the above equation is very difficult to solve (numerically) for a given  $y_0$ , but easy to solve for a value  $\tilde{y}$  "near"  $y_0$ . Then, if  $T^{-1}$  is continuous, the correspondent solutions  $x_0$  and  $\tilde{x}$  are also "near" since

$$\|x_0 - \tilde{x}\| = \|T^{-1}y_0 - T^{-1}\tilde{y}\| \leq \|T^{-1}\|\|y_0 - \tilde{y}\|$$

Therefore we can solve the equation for a "near" value  $\tilde{y}$  instead, without obtaining a significant error.