



matrix inversion lemma

| | |
|------------------|--------------------------|
| Canonical name | MatrixInversionLemma |
| Date of creation | 2013-03-22 15:38:44 |
| Last modified on | 2013-03-22 15:38:44 |
| Owner | kronos (12218) |
| Last modified by | kronos (12218) |
| Numerical id | 6 |
| Author | kronos (12218) |
| Entry type | Result |
| Classification | msc 47S99 |
| Synonym | Sherman-Morrison formula |
| Synonym | Woodbury matrix identity |
| Synonym | Woodbury formula |
| Synonym | rank-k correction |
| Related topic | SchurComplement |

These frequently used formulae allow to quickly calculate the inverse of a slight modification of an operator (matrix) x , given that x^{-1} is already known.

The matrix inversion lemma states that

$$(x + s\sigma z^*)^{-1} = x^{-1} - x^{-1}s(\sigma^{-1} + z^*x^{-1}s)^{-1}z^*x^{-1},$$

where x , s , z^* and σ are operators (matrices) of appropriate size. The formula especially is convenient if the rank of the regular σ is 1, or small in comparison to x 's rank.

This identity, involving the inverse of Schur's complement $d - z^*x^{-1}s$ (hopefully this may be easily computed) holds as well:

$$\begin{bmatrix} x & s \\ z^* & d \end{bmatrix}^{-1} = \begin{bmatrix} x^{-1} + x^{-1}s(d - z^*x^{-1}s)^{-1}z^*x^{-1} & -x^{-1}s(d - z^*x^{-1}s)^{-1} \\ -(d - z^*x^{-1}s)^{-1}z^*x^{-1} & (d - z^*x^{-1}s)^{-1} \end{bmatrix}.$$