



Math for the people, by the people.

ergodicity of a map in terms of its induced operator

Canonical name	ErgodicityOfAMapInTermsOfItsInducedOperator
Date of creation	2013-03-22 17:59:22
Last modified on	2013-03-22 17:59:22
Owner	asteroid (17536)
Last modified by	asteroid (17536)
Numerical id	6
Author	asteroid (17536)
Entry type	Theorem
Classification	msc 47A35
Classification	msc 37A30
Classification	msc 37A25
Classification	msc 28D05

Theorem - Let (X, \mathfrak{B}, μ) be a probability space and $T : X \longrightarrow X$ a measure-preserving transformation. The following statements are equivalent:

1. - T is ergodic.
2. - If f is a measurable function and $f \circ T = f$ a.e., then f is constant a.e.
3. - If f is a measurable function and $f \circ T \geq f$ a.e., then f is constant a.e.
4. - If $f \in L^2(X)$ and $f \circ T = f$ a.e., then f is constant a.e..
5. - If $f \in L^p(X)$, with $p \geq 1$, and $f \circ T = f$ a.e., then f is constant a.e.

Let U_T denote the operator induced by T , i.e. the operator defined by $U_T f := f \circ T$. The statements above are statements about U_T . The above theorem can be rewritten as follows:

Theorem - Let (X, \mathfrak{B}, μ) be a probability space and $T : X \longrightarrow X$ a measure-preserving transformation. The following statements are equivalent:

1. - T is ergodic.
2. - The only fixed points of U_T are the functions that are constant a.e.
3. - If f a measurable function and $U_T f \geq f$ a.e., then f is constant a.e.
4. - The eigenspace of U_T (seen as an operator in $L^p(X)$, with $p \geq 1$) associated with the eigenvalue 1, is one-dimensional and consists of functions that are constant a.e.