



Lagrange multiplier method

Canonical name	LagrangeMultiplierMethod
Date of creation	2013-03-22 12:25:10
Last modified on	2013-03-22 12:25:10
Owner	cvalente (11260)
Last modified by	cvalente (11260)
Numerical id	10
Author	cvalente (11260)
Entry type	Definition
Classification	msc 49K30
Related topic	ExampleOfCalculusOfVariations
Related topic	IsoperimetricProblem
Defines	Lagrange multiplier

The Lagrange multiplier method is used when one needs to find the extreme or stationary points of a function on a set which is a subset of the domain.

Method

Suppose that $f(\mathbf{x})$ and $g_i(\mathbf{x}), i = 1, \dots, m$ ($\mathbf{x} \in \mathbb{R}^n$) are differentiable functions that map $\mathbb{R}^n \mapsto \mathbb{R}$, and we want to solve

$$\min f(\mathbf{x}), \max f(\mathbf{x}) \quad \text{such that} \quad g_i(\mathbf{x}) = 0, \quad i = 1, \dots, m$$

By a calculus theorem, if the constraints are independent, the gradient of f , ∇f , must satisfy the following equation at the stationary points:

$$\nabla f = \sum_{i=1}^m \lambda_i \nabla g_i$$

The constraints are said to be independent iff all the gradients of each constraint are linearly independent, that is:

$\{\nabla g_1(\mathbf{x}), \dots, \nabla g_m(\mathbf{x})\}$ is a set of linearly independent vectors on all points where the constraints are verified.

Note that this is equivalent to finding the stationary points of:

$$f(\mathbf{x}) - \sum_{i=1}^m \lambda_i (g_i(\mathbf{x}))$$

for \mathbf{x} in the domain and the *Lagrange multipliers* λ_i without restrictions.

After finding those points, one applies the g_i constraints to get the actual stationary points subject to the constraints.