



planetmath.org

Math for the people, by the people.

Lagrange multipliers on Banach spaces

Canonical name	LagrangeMultipliersOnBanachSpaces
Date of creation	2013-03-22 15:28:30
Last modified on	2013-03-22 15:28:30
Owner	stevecheng (10074)
Last modified by	stevecheng (10074)
Numerical id	5
Author	stevecheng (10074)
Entry type	Theorem
Classification	msc 49-00
Classification	msc 49K35

Let U be open in a real Banach space X , and Y be another real Banach space. Let $f: U \rightarrow \mathbb{R}$ and $g: U \rightarrow Y$ be continuously differentiable functions.

Suppose that a is a minimum or maximum point of f on $M = \{x \in U : g(x) = 0\}$, and the Fréchet derivative $Dg(a): X \rightarrow Y$ is surjective. Then there exists a Lagrange multiplier vector $\lambda \in Y^*$ such that

$$Df(a) = Dg(a)^*\lambda = \lambda \circ Dg(a).$$

(The function $Dg(a)^*: Y^* \rightarrow X^*$ denotes the pullback or adjoint by $Dg(a)$ on the continuous duals, defined by the second equality.)

If X and Y are finite-dimensional, writing out the above equation in matrix form shows that λ really is the usual Lagrange multiplier vector. The condition that $Dg(a)$ is surjective means that $Dg(a)$ must have full rank as a matrix.

References

- [1] Eberhard Zeidler. *Applied functional analysis: main principles and their applications*. Springer-Verlag, 1995.