



planetmath.org

Math for the people, by the people.

example of using Lagrange multipliers

Canonical name	ExampleOfUsingLagrangeMultipliers
Date of creation	2013-03-22 18:48:12
Last modified on	2013-03-22 18:48:12
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	7
Author	pahio (2872)
Entry type	Example
Classification	msc 51N20
Classification	msc 26B10
Synonym	example of Lagrange multipliers
Related topic	ParallelismOfTwoPlanes
Related topic	ExampleNeedingTwoLagrangeMultipliers

One to determine the perpendicular distance of the parallel planes

$$Ax + By + Cz + D = 0 \quad \text{and} \quad Ax + By + Cz + E = 0$$

is to use the Lagrange multiplier method. In this case we may to minimise the Euclidean distance of a point  $(x, y, z)$  of the former plane to a (fixed) point  $(x_0, y_0, z_0)$  of the latter plane.

Thus we have the equation  $Ax_0 + By_0 + Cz_0 + E = 0$  which we can subtract from the first plane equation, getting

$$g := A(x - x_0) + B(y - y_0) + C(z - x_0) + D - E = 0. \quad (1)$$

This is the (only) constraint equation for minimising the <http://planetmath.org/SquareOfANumber>

$$f := (x - x_0)^2 + (y - y_0)^2 + (z - x_0)^2 \quad (2)$$

of the distance of the points.

The polynomial functions  $f$  and  $g$  satisfy the differentiability requirements. Accordingly, we can find the minimising point  $(x, y, z)$  by considering the system of equations formed by (1) and

$$\begin{cases} \frac{\partial f}{\partial x} + \lambda \frac{\partial g}{\partial x} \equiv 2(x - x_0) + \lambda A = 0, \\ \frac{\partial f}{\partial y} + \lambda \frac{\partial g}{\partial y} \equiv 2(y - y_0) + \lambda B = 0, \\ \frac{\partial f}{\partial z} + \lambda \frac{\partial g}{\partial z} \equiv 2(z - z_0) + \lambda C = 0. \end{cases} \quad (3)$$

We solve from (3) the differences

$$x - x_0 = -\frac{A\lambda}{2}, \quad y - y_0 = -\frac{B\lambda}{2}, \quad z - z_0 = -\frac{C\lambda}{2}$$

and set them into (1). It then yields the value

$$\lambda = \frac{2(D - E)}{A^2 + B^2 + C^2}$$

of the Lagrange multiplier, which we substitute into the preceding three equations obtaining

$$x - x_0 = \frac{A(D - E)}{A^2 + B^2 + C^2}, \quad y - y_0 = \frac{B(D - E)}{A^2 + B^2 + C^2}, \quad z - z_0 = \frac{C(D - E)}{A^2 + B^2 + C^2}.$$

These values give the minimal distance when put into the expression of  $\sqrt{f}$ :

$$d = \sqrt{\frac{(D - E)^2(A^2 + B^2 + C^2)}{(A^2 + B^2 + C^2)^2}}.$$

Hence we have gotten the distance

$$d = \frac{|D - E|}{\sqrt{A^2 + B^2 + C^2}}.$$