



planetmath.org

Math for the people, by the people.

Szemerédi-Trotter theorem

Canonical name	SzemerediTrotterTheorem
Date of creation	2013-03-22 13:21:30
Last modified on	2013-03-22 13:21:30
Owner	bbukh (348)
Last modified by	bbukh (348)
Numerical id	7
Author	bbukh (348)
Entry type	Theorem
Classification	msc 51A20
Classification	msc 05C10

The number of incidences of a set of n points and a set of m lines in the real plane \mathbb{R}^2 is

$$I = O(n + m + (nm)^{\frac{2}{3}}).$$

Proof. Let's consider the points as vertices of a graph, and connect two vertices by an edge if they are adjacent on some line. Then the number of edges is $e = I - m$. If $e < 4n$ then we are done. If $e \geq 4n$ then by crossing lemma

$$m^2 \geq \text{cr}(G) \geq \frac{1}{64} \frac{(I - m)^3}{n^2},$$

and the theorem follows.

Recently, Tóth[?] extended the theorem to the complex plane \mathbb{C}^2 . The proof is difficult.

References

- [1] Csaba D. Tóth. The Szemerédi-Trotter theorem in the complex plane.
<http://www.arxiv.org/abs/math.CO/0305283>arXiv:CO/0305283,
 May 2003.