



planetmath.org

Math for the people, by the people.

volume as integral

Canonical name	VolumeAsIntegral
Date of creation	2013-03-22 17:20:44
Last modified on	2013-03-22 17:20:44
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	9
Author	pahio (2872)
Entry type	Topic
Classification	msc 51M25
Classification	msc 51-00
Related topic	Volume
Related topic	VolumeOfSolidOfRevolution
Related topic	RiemannMultipleIntegral
Related topic	ExampleOfRiemannTripleIntegral

The <http://planetmath.org/VolumeOfSolidOfRevolution> volume of a solid of revolution can be obtained from

$$V = \int_a^b \pi[f(x)]^2 dx,$$

where the integrand is the area of the intersection disc of the solid of revolution and a plane perpendicular to the axis of revolution at a certain value of  $x$ . This volume formula may be generalized to an analogous formula containing instead of the area  $\pi[f(x)]^2$  a more general intersection area  $A(t)$  obtained from a given solid by cutting it with a set of parallel planes determined by the parameter  $t$  on a certain axis. One must assume that the function  $t \mapsto A(t)$  is continuous on an interval  $[a, b]$  where  $a$  and  $b$  correspond to the “ends” of the solid. If the  $t$ -axis <http://planetmath.org/AngleBetweenTwoLines> forms an angle  $\omega$  with the normal line of those planes, then we have the volume formula of the form

$$V = \int_a^b A(t) dt \cos \omega.$$