

planetmath.org

Math for the people, by the people.

polarities and forms

Canonical name Polarities And Forms
Date of creation 2013-03-22 15:58:13
Last modified on 2013-03-22 15:58:13
Owner Algeboy (12884)
Last modified by Algeboy (12884)

Numerical id 7

Author Algeboy (12884)

Entry type Topic

Classification msc 51A05 Related topic polarity Related topic Projectivity

Related topic ProjectiveGeometry

Related topic Isometry2

Related topic ProjectiveGeometry3
Related topic ClassicalGroups

Related topic Polarity2

Related topic DualityWithRespectToANonDegenerateBilinearForm

Through out this article we assume $\dim V \neq 2$. This is not a true constraint as there are only trivial dualities for $\dim V \leq 2$.

Proposition 1. Every duality gives rise to a non-degenerate sesquilinear form, and visa-versa.

Proof. To see this, let $d: PG(V) \to PG(V)$ be a duality. We may express this as an order preserving map $d: PG(V) \to PG(V^*)$. Then by the fundamental theorem of projective geometry it follows d is induced by a bijective semi-linear transformation $\hat{d}: V \to V^*$.

An semi-linear isomorphism of V to V^* is equivalent to specifying a non-degenerate sesquilinear form. In particular, define the form $b: V \times V \to k$ by $b(v, w) = (v)(w\hat{d})$ (notice $w\hat{d} \in V^*$ so $w\hat{d}: V \to k$).

Now, if $b: V \times V \to k$ is a non-degenerate sesquilinear form. Then define

$$\hat{b}: V \to V^*: v \mapsto b(-,v): V \to k$$

which is semi-linear, as b is sesquilinear, and bijective, since b is non-degenerate. Therefore \hat{b} induces an order preserving bijection $PG(V) \to PG(V^*)$, that is, a duality.

We write W^{\perp} for the image of the induced duality of a non-degenerate sesquilinear form b. Notice that $W^{\perp} = \{w \in V : b(v, W) = 0\}$. (Although the form may not be reflexive, we still use the \perp notation, but we now demonstrate that we can indeed specialize to the reflexive case.) Notice then that

$$\dim W^{\perp} = \dim V - \dim W$$
.

Corollary 2. Every polarity gives rise to a reflexive non-degenerate sesquilinear form, and visa-versa.

Proof. Let b be the sesquilinear form induced by the polarity p. Then suppose we have $v, w \in V$ such that $0 = b(v, w) = (v)(w\hat{p})$. So $\langle v \rangle \leq \langle w \rangle^{\perp} = \langle w \rangle p$. But p has order 2 so $\langle v \rangle^{\perp} = \langle v \rangle p \geq \langle w \rangle$. But this implies b(w, v) = 0 so b is reflexive.

Likewise, given a reflexive non-degenerate sesquilinear form b it gives rise do a duality d induced by \hat{b} . By the reflexivity, $b(W, W^{\perp}) = 0$ implies $b(W^{\perp}, W) = 0$ also. As $(W^{\perp})^{\perp} = \{v \in V : b(v, (W^{\perp})^{\perp}) = 0\}$ it follows $W \leq (W^{\perp})^{\perp}$. But by dimension arguments:

$$\dim(W^{\perp})^{\perp} = \dim V - \dim W^{\perp} = \dim V - (\dim V - \dim W) = \dim W$$
 we conclude $W = (W^{\perp})^{\perp}$. Thus d is a polarity. \square

From the fundamental theorem of projective geometry it follows if dim $V \neq 2$ then every order preserving map is induced by a semi-linear transformation of V. In similar fashion we have

Proposition 3. $P\Gamma L^*(V) = P\Gamma L(V) \rtimes \mathbb{Z}_2$, meaning that every order reversing map $f: PG(V) \to PG(V)$ can be decomposed as a f = sr where s is induced from a semi-linear transformation and r is a polarity.

Proof. Let d be any duality of PG(V). Then d^2 is order preserving. Thus d^2 is a projectivity so by the fundamental theorem of projective geometry d^2 is induced by a semi-linear transformation s. Therefore $P\Gamma L(V)$ has index 2 in $P\Gamma L^*(V)$. Finally it suffices to provide any polarity of PG(V) to prove $P\Gamma L^*(V) = P\Gamma L(V) \rtimes \mathbb{Z}_2$. For this use any reflexive non-degenerate sesquilinear form.

Remark 4. The group $P\Gamma L^*(V)$ is the automorphism group of PSL(V). In particular, the polarities account for the graph automorphisms of the Dynkin diagram of A_{d-1} , $d = \dim V$. When $\dim V = 2$ there is no graph automorphism, just as there are no dualities (points are hyperplanes when $\dim V = 2$.)

References

- [1] Gruenberg, K. W. and Weir, A.J. *Linear Geometry 2nd Ed.* (English) [B] Graduate Texts in Mathematics. 49. New York Heidelberg Berlin: Springer-Verlag. X, 198 p. DM 29.10; \$ 12.80 (1977).
- [2] Kantor, W. M. Lectures notes on Classical Groups.