



**planetmath.org**

Math for the people, by the people.

## proof of tangents law

|                  |                     |
|------------------|---------------------|
| Canonical name   | ProofOfTangentsLaw  |
| Date of creation | 2013-03-22 13:11:04 |
| Last modified on | 2013-03-22 13:11:04 |
| Owner            | CWoo (3771)         |
| Last modified by | CWoo (3771)         |
| Numerical id     | 6                   |
| Author           | CWoo (3771)         |
| Entry type       | Proof               |
| Classification   | msc 51-00           |

To prove that

$$\frac{a-b}{a+b} = \frac{\tan(\frac{A-B}{2})}{\tan(\frac{A+B}{2})}$$

we start with the sines law, which says that

$$\frac{a}{\sin(A)} = \frac{b}{\sin(B)}.$$

This implies that

$$a \sin(B) = b \sin(A)$$

We can write  $\sin(A)$  as

$$\sin(A) = \sin(\frac{A+B}{2}) \cos(\frac{A-B}{2}) + \cos(\frac{A+B}{2}) \sin(\frac{A-B}{2}).$$

and  $\sin(B)$  as

$$\sin(B) = \sin(\frac{A+B}{2}) \cos(\frac{A-B}{2}) - \cos(\frac{A+B}{2}) \sin(\frac{A-B}{2}).$$

Therefore, we have

$$a(\sin(\frac{A+B}{2}) \cos(\frac{A-B}{2}) - \cos(\frac{A+B}{2}) \sin(\frac{A-B}{2})) = b(\sin(\frac{A+B}{2}) \cos(\frac{A-B}{2}) + \cos(\frac{A+B}{2}) \sin(\frac{A-B}{2}))$$

Dividing both sides by  $\cos(\frac{A-B}{2}) \cos(\frac{A+B}{2})$ , we have,

$$a(\tan(\frac{A+B}{2}) - \tan(\frac{A-B}{2})) = b(\tan(\frac{A+B}{2}) + \tan(\frac{A-B}{2}))$$

This gives us

$$\frac{a}{b} = \frac{\tan(\frac{A+B}{2}) + \tan(\frac{A-B}{2})}{\tan(\frac{A+B}{2}) - \tan(\frac{A-B}{2})}$$

Hence we find that

$$\frac{a-b}{a+b} = \frac{\frac{a}{b} - 1}{\frac{a}{b} + 1} = \frac{\tan(\frac{A-B}{2})}{\tan(\frac{A+B}{2})}.$$