



planetmath.org

Math for the people, by the people.

isogonal trajectory

Canonical name	IsogonalTrajectory
Date of creation	2013-03-22 18:59:20
Last modified on	2013-03-22 18:59:20
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	7
Author	pahio (2872)
Entry type	Derivation
Classification	msc 51N20
Classification	msc 34A26
Classification	msc 34A09
Related topic	AngleBetweenTwoCurves
Related topic	OrthogonalCurves
Related topic	ExampleOfIsogonalTrajectory
Related topic	AngleBetweenTwoLines
Defines	isogonal trajectory

Let a one-parametric family of plane curves γ have the differential equation

$$F(x, y, \frac{dy}{dx}) = 0. \quad (1)$$

We want to determine the *isogonal trajectories* of this family, i.e. the curves ι intersecting all members of the family under a given angle, which is denoted by ω . For this purpose, we denote the slope angle of any curve γ at such an intersection point by α and the slope angle of ι at the same point by β . Then

$$\beta - \alpha = \omega \quad (\text{or alternatively } -\omega),$$

and accordingly

$$\frac{dy}{dx} = \tan \alpha = \frac{\tan \beta - \tan \omega}{1 + \tan \beta \tan \omega} = \frac{y' - \tan \omega}{1 + y' \tan \omega},$$

where y' means the slope of ι . Thus the equation

$$F(x, y, \frac{y' - \tan \omega}{1 + y' \tan \omega}) = 0 \quad (2)$$

is satisfied by the derivative y' of the ordinate of ι . In other , (2) is the differential equation of all isogonal trajectories of the given family of curves.

Note. In the special case $\omega = \frac{\pi}{2}$, it's a question of orthogonal trajectories.