



planetmath.org

Math for the people, by the people.

affine combination

Canonical name	AffineCombination
Date of creation	2013-03-22 16:00:13
Last modified on	2013-03-22 16:00:13
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	19
Author	CWoo (3771)
Entry type	Definition
Classification	msc 51A15
Synonym	affine independent
Related topic	AffineGeometry
Related topic	AffineTransformation
Related topic	ConvexCombination
Defines	affine independence
Defines	affinely independent
Defines	affine hull

Definition

Let V be a vector space over a division ring D . An *affine combination* of a finite set of vectors $v_1, \dots, v_n \in V$ is a linear combination of the vectors

$$k_1 v_1 + \dots + k_n v_n$$

such that $k_i \in D$ subject to the condition $k_1 + \dots + k_n = 1$. In effect, an affine combination is a weighted average of the vectors in question.

For example, $v = \frac{1}{2}v_1 + \frac{1}{2}v_2$ is an affine combination of v_1 and v_2 provided that the characteristic of D is not 2. v is known as the midpoint of v_1 and v_2 . More generally, if $\text{char}(D)$ does not divide m , then

$$v = \frac{1}{m}(v_1 + \dots + v_m)$$

is an affine combination of the v_i 's. v is the barycenter of v_1, \dots, v_n .

Relations with Affine Subspaces

Assume now $\text{char}(D) = 0$. Given $v_1, \dots, v_n \in V$, we can form the set A of all affine combinations of the v_i 's. We have the following

A is a finite dimensional affine subspace. Conversely, a finite dimensional affine subspace A is the set of all affine combinations of a finite set of vectors in A .

Proof. Suppose A is the set of affine combinations of v_1, \dots, v_n . If $n = 1$, then A is a singleton $\{v\}$, so $A = 0 + v$, where 0 is the null subspace of V . If $n > 1$, we may pick a non-zero vector $v \in A$. Define $S = \{a - v \mid a \in A\}$. Then for any $s \in S$ and $d \in D$, $ds = d(a - v) = da + (1 - d)v - v$. Since $da + (1 - d)v \in A$, $ds \in S$. If $s_1, s_2 \in S$, then $\frac{1}{2}(s_1 + s_2) = \frac{1}{2}((a_1 - v) + (a_2 - v)) = \frac{1}{2}(a_1 + a_2) - v \in S$, since $\frac{1}{2}(a_1 + a_2) \in A$. So $\frac{1}{2}(s_1 + s_2) \in S$. Therefore, $s_1 + s_2 = 2(\frac{1}{2}(s_1 + s_2)) \in S$. This shows that S is a vector subspace of V and that $A = S + v$ is an affine subspace.

Conversely, let A be a finite dimensional affine subspace. Write $A = S + v$, where S is a subspace of V . Since $\dim(S) = \dim(A) = n$, S has a basis $\{s_1, \dots, s_n\}$. For each $i = 1, \dots, n$, define $v_i = ns_i + v$. Given $a \in A$, we

have

$$\begin{aligned}
a &= s + v = k_1 s_1 + \cdots + k_n s_n + v \\
&= \frac{k_1}{n}(v_1 - v) + \cdots + \frac{k_n}{n}(v_n - v) + v \\
&= \frac{k_1}{n}v_1 + \cdots + \frac{k_n}{n}v_n + \left(1 - \frac{k_1}{n} - \cdots - \frac{k_n}{n}\right)v.
\end{aligned}$$

From this calculation, it is evident that a is an affine combination of v_1, \dots, v_n , and v . \square

When A is the set of affine combinations of two distinct vectors v, w , we see that A is a line, in the sense that $A = S + v$, a translate of a one-dimensional subspace S (a line through 0). Every element in A has the form $dv + (1 - d)w$, $d \in D$. Inspecting the first part of the proof in the previous proposition, we see that the argument involves no more than two vectors at a time, so the following useful corollary is apparant:

A is an affine subspace iff for every pair of vectors in A , the line formed by the pair is also in A .

Note, however, that the A in the above corollary is not assumed to be finite dimensional.

Remarks.

- If one of v_1, \dots, v_n is the zero vector, then A coincides with S . In other words, an affine subspace is a vector subspace if it contains the zero vector.
- Given $A = \{k_1 v_1 + \cdots + k_n v_n \mid v_i \in V, k_i \in D, \sum k_i = 1\}$, the subset

$$\{k_1 v_1 + \cdots + k_n v_n \in A \mid k_i = 0\}$$

is also an affine subspace.

Affine Independence

Since every element in a finite dimensional affine subspace A is an affine combination of a finite set of vectors in A , we have the similar concept of a spanning set of an affine subspace. A minimal spanning set M of an affine subspace is said to be *affinely independent*. We have the following three equivalent characterization of an affinely independent subset M of a finite dimensional affine subspace:

1. $M = \{v_1, \dots, v_n\}$ is affinely independent.
2. every element in A can be written as an affine combination of elements in M in a *unique* fashion.
3. for every $v \in M$, $N = \{v_i - v \mid v \neq v_i\}$ is linearly independent.

Proof. We will proceed as follows: (1) implies (2) implies (3) implies (1).

(1) implies (2). If $a \in A$ has two distinct representations $k_1v_1 + \dots + k_nv_n = a = r_1v_1 + \dots + r_nv_n$, we may assume, say $k_1 \neq r_1$. So $k_1 - r_1$ is invertible with inverse $t \in D$. Then

$$v_1 = t(r_2 - k_2)v_2 + \dots + t(r_n - k_n)v_n.$$

Furthermore,

$$\sum_{i=2}^n t(r_i - k_i) = t\left(\sum_{i=2}^n r_i - \sum_{i=2}^n k_i\right) = t(1 - r_1 - 1 + k_1) = 1.$$

So for any $b \in A$, we have

$$b = s_1v_1 + \dots + s_nv_n = s_1(t(r_2 - k_2)v_2 + \dots + t(r_n - k_n)v_n) + \dots + s_nv_n.$$

The sum of the coefficients is easily seen to be 1, which implies that $\{v_2, \dots, v_n\}$ is a spanning set of A that is smaller than M , a contradiction.

(2) implies (3). Pick $v = v_1$. Suppose $0 = s_2(v_2 - v_1) + \dots + s_n(v_n - v_1)$. Expand and we have $0 = (-s_2 - \dots - s_n)v_1 + s_2v_2 + \dots + s_nv_n$. So $(1 - s_2 - \dots - s_n)v_1 + s_2v_2 + \dots + s_nv_n = v_1 \in A$. By assumption, there is exactly one way to express v_1 , so we conclude that $s_2 = \dots = s_n = 0$.

(3) implies (1). If M were not minimal, then some $v \in M$ could be expressed as an affine combination of the remaining vectors in M . So suppose $v_1 = k_2v_2 + \dots + k_nv_n$. Since $\sum k_i = 1$, we can rewrite this as $0 = k_2(v_2 - v_1) + \dots + k_n(v_n - v_1)$. Since not all $k_i = 0$, $N = \{v_2 - v_1, \dots, v_n - v_1\}$ is not linearly independent. \square

Remarks.

- If $\{v_1, \dots, v_n\}$ is affinely independent set spanning A , then $\dim(A) = n - 1$.

- More generally, a set M (not necessarily finite) of vectors is said to be affinely independent if there is a vector $v \in M$, such that $N = \{w - v \mid v \neq w \in M\}$ is linearly independent (every finite subset of N is linearly independent). The above three characterizations are still valid in this general setting. However, one must be careful that an affine combination is a finitary operation so that when we take the sum of an infinite number of vectors, we have to realize that only a finite number of them are non-zero.
- Given any set S of vectors, the *affine hull* of S is the smallest affine subspace A that contains every vector of S , denoted by $\text{Aff}(S)$. Every vector in $\text{Aff}(S)$ can be written as an affine combination of vectors in S .