



convex functions lie above their supporting lines

Canonical name	ConvexFunctionsLieAboveTheirSupportingLines
Date of creation	2013-03-22 16:59:20
Last modified on	2013-03-22 16:59:20
Owner	Andrea Ambrosio (7332)
Last modified by	Andrea Ambrosio (7332)
Numerical id	5
Author	Andrea Ambrosio (7332)
Entry type	Result
Classification	msc 52A41
Classification	msc 26A51
Classification	msc 26B25

Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be a convex, twice differentiable function on $[a, b]$. Then $f(x)$ lies above its supporting lines, i.e. it's greater than any tangent line in $[a, b]$.

Proof. :

Let $r(x) = f(x_0) + f'(x_0)(x - x_0)$ be the tangent of $f(x)$ in $x = x_0 \in [a, b]$.

By Taylor theorem, with remainder in Lagrange form, one has, for any $x \in [a, b]$:

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{1}{2}f''(\xi(x))(x - x_0)^2$$

with $\xi(x) \in [a, b]$. Then

$$f(x) - r(x) = \frac{1}{2}f''(\xi(x))(x - x_0)^2 \geq 0$$

since $f''(\xi(x)) \geq 0$ by convexity. □