



Math for the people, by the people.

germ space

| | |
|------------------|---------------------|
| Canonical name | GermSpace |
| Date of creation | 2013-03-22 19:18:20 |
| Last modified on | 2013-03-22 19:18:20 |
| Owner | joking (16130) |
| Last modified by | joking (16130) |
| Numerical id | 4 |
| Author | joking (16130) |
| Entry type | Definition |
| Classification | msc 53B99 |

Let X, Y be topological spaces and $x \in X$. Consider the set of all continuous functions

$$C(X, Y) = \{f : X \rightarrow Y \mid f \text{ is continuous}\}.$$

For any two functions $f, g : X \rightarrow Y$ we put

$$f \sim_x g$$

if and only if there exists an open neighbourhood $U \subseteq X$ of x such that

$$f|_U = g|_U.$$

The corresponding quotient set is called **the germ space** at $x \in X$ and we denote it by $G_x(X, Y)$.

More generally, if X, Y are topological spaces with $x \in X$, then consider the following set:

$$C_x(X, Y) = \{f : U \rightarrow Y \mid f \text{ is continuous and } U \text{ is an open neighbourhood of } x\}.$$

Again we define a relation on $C_x(X, Y)$. If $f : U \rightarrow Y$ and $g : U' \rightarrow Y$, then put

$$f \sim_x g$$

if and only if there exists an open neighbourhood $V \subseteq X$ of x such that $V \subseteq U \cap U'$ and

$$f|_V = g|_V.$$

The corresponding set is called **the generalized germ space** at $x \in X$ and we denote it by $G_x^*(X, Y)$.

Note that if $Y = \mathbb{R}$ or $Y = \mathbb{C}$ (or Y is any topological ring), then both $G_x(X, Y)$ and $G_x^*(X, Y)$ have a well-defined ring structure via pointwise addition and multiplication.