



Math for the people, by the people.

fundamental theorem of space curves

Canonical name	FundamentalTheoremOfSpaceCurves
Date of creation	2013-03-22 13:23:28
Last modified on	2013-03-22 13:23:28
Owner	rmilson (146)
Last modified by	rmilson (146)
Numerical id	5
Author	rmilson (146)
Entry type	Theorem
Classification	msc 53A04
Related topic	SpaceCurve

Informal summary. The curvature and torsion of a space curve are invariant with respect to Euclidean motions. Conversely, a given space curve is determined up to a Euclidean motion, by its curvature and torsion, expressed as functions of the arclength.

Theorem. Let $\gamma : I \rightarrow \mathbb{R}^3$ be a regular, parameterized space curve, without points of inflection. Let $\kappa(t), \tau(t)$ be the corresponding curvature and torsion functions. Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a Euclidean isometry. The curvature and torsion of the transformed curve $T(\gamma(t))$ are given by $\kappa(t)$ and $\tau(t)$, respectively.

Conversely, let $\kappa, \tau : I \rightarrow \mathbb{R}$ be continuous functions, defined on an interval $I \subset \mathbb{R}$, and suppose that $\kappa(t)$ never vanishes. Then, there exists an arclength parameterization $\gamma : I \rightarrow \mathbb{R}^3$ of a regular, oriented space curve, without points of inflection, such that $\kappa(t)$ and $\tau(t)$ are the corresponding curvature and torsion functions. If $\hat{\gamma} : I \rightarrow \mathbb{R}^3$ is another such space curve, then there exists a Euclidean isometry $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ such that $\hat{\gamma}(t) = T(\gamma(t))$.