



planetmath.org

Math for the people, by the people.

proof of Poincaré lemma

Canonical name	ProofOfPoincareLemma
Date of creation	2013-03-22 14:24:36
Last modified on	2013-03-22 14:24:36
Owner	pbruin (1001)
Last modified by	pbruin (1001)
Numerical id	4
Author	pbruin (1001)
Entry type	Proof
Classification	msc 53-00
Classification	msc 55N05

Let X be a smooth manifold, and let ω be a closed differential form of degree $k > 0$ on X . For any $x \in X$, there exists a contractible neighbourhood $U \subset X$ of x (i.e. U is homotopy equivalent to a single point), with inclusion map

$$\iota: U \hookrightarrow X.$$

To construct such a neighbourhood, take for example an open ball in a coordinate chart around x . Because of the homotopy invariance of de Rham cohomology, the k th de Rham cohomology group $H^k(U)$ is isomorphic to that of a point; in particular,

$$H^k(U) = 0 \quad \text{for all } k > 0.$$

Since $d(\iota^*\omega) = \iota^*(d\omega) = 0$, this implies that there exists a $(k-1)$ -form η on U such that $d\eta = \iota^*\omega$. In the case where X is a contractible manifold, such an η exists globally since we can choose $U = X$ above.