



**planetmath.org**

Math for the people, by the people.

## every symplectic manifold has even dimension

Canonical name	EverySymplecticManifoldHasEvenDimension
Date of creation	2013-03-22 15:44:05
Last modified on	2013-03-22 15:44:05
Owner	cvalente (11260)
Last modified by	cvalente (11260)
Numerical id	18
Author	cvalente (11260)
Entry type	Theorem
Classification	msc 53D05
Related topic	AlternatingForm

All we need to prove is that every finite dimensional vector space  $V$  with an anti-symmetric non-degenerate linear form  $\omega$  has an even dimension  $2k$ . This is only a linear algebra result. In the case of a symplectic manifold  $V$  is just the tangent space at a point, and thus its dimension equals the manifold's dimension.

Pick any not null vector  $v_0 \in V$ . Since  $\omega$  is non-degenerate  $\omega(v_0, \cdot)$  is a not null linear form. Therefore there exists a not null vector  $u_0$  such that  $\omega(v_0, u_0) = 1$

Now  $v_0$  and  $u_0$  are linearly independent because if  $v_0 = \lambda u_0$  then  $\omega(v_0, u_0) = \omega(\lambda u_0, u_0) = \lambda \omega(u_0, u_0) = 0$  (by anti-symmetry).

Let  $V_0 = \text{span}\{v_0, u_0\}$ . Consider a space  $V_1$  of "orthogonal" elements to  $V_0$  under  $\omega$ . That is:

$$V_1 = \{v_1 \in V : \text{for all } v \in V_0, \omega(v, v_1) = 0\}$$

We now prove  $V = V_0 \oplus V_1$ :

- $V_0 \cap V_1 = \{0\}$

Suppose  $w \in V_0 \cap V_1$  is not null, then it can be written  $w = \alpha v_0 + \beta u_0$  because it belongs to  $V_0$ . Since it also belongs to  $V_1$  is is "orthogonal" to both  $v_0$  and  $u_0$ . That is:

$$\omega(v_0, w) = 0 \implies \beta \omega(v_0, u_0) = 0 \implies \beta = 0 \text{ similarly}$$

$$\omega(u_0, w) = 0 \implies \alpha \omega(u_0, v_0) = 0 \implies \alpha = 0$$

So  $w$  must be null.

- $V = V_0 \oplus V_1$

Suppose  $w \in V$ . Let  $\alpha = \omega(v_0, w)$ ,  $\beta = \omega(u_0, w)$ ,  $w_0 = \alpha u_0 - \beta v_0$ .

Then  $\omega(v_0, w_0) = \alpha = \omega(v_0, w)$  and  $\omega(u_0, w_0) = \beta = \omega(u_0, w)$ .

Considering  $w_1 = w - w_0$  we have  $w = w_0 + w_1$  (by construction) and  $\omega(v_0, w_1) = \omega(v_0, w - w_0) = \omega(v_0, w) - \omega(v_0, w_0) = \omega(v_0, w) - \omega(v_0, w) = 0$  and similarly for  $\omega(u_0, w_1)$

So  $w_1 \in V_1$ ,  $w_0 \in V_0$  and  $w = w_0 + w_1$  and thus  $V = V_0 \oplus V_1$

So the matrix representation of  $\omega$  is block-diagonal in  $V_0 \oplus V_1$  and a restriction anti-symmetric bilinear for of  $\omega$  to  $V_1$  exists.

If  $V_1$  is not null we can repeat the procedure with the restriction. Since  $\dim(V) = \dim(V_0) + \dim(V_1)$  and  $V$  is finite dimensional the procedure must stop at a finite step.

At the end we get a decomposition  $V = \bigoplus_{i=0}^{k-1} V_i$ , where  $\dim(V_i) = 2$  and  $\dim(V) = 2k$  is even.