



Math for the people, by the people.

Nash isometric embedding theorem

Canonical name	NashIsometricEmbeddingTheorem
Date of creation	2013-03-22 15:38:17
Last modified on	2013-03-22 15:38:17
Owner	Simone (5904)
Last modified by	Simone (5904)
Numerical id	7
Author	Simone (5904)
Entry type	Theorem
Classification	msc 53C20
Classification	msc 53C42
Classification	msc 57R40
Classification	msc 58A05

Every compact n -dimensional Riemannian manifold M of class C^k ($3 \leq k \leq \infty$) can be C^k -isometrically imbedded in any small portion of a Euclidean space \mathbb{R}^N , where

$$N = \frac{1}{2}n(3n + 11).$$

Every non-compact n -dimensional Riemannian manifold M of class C^k ($3 \leq k \leq \infty$) can be C^k -isometrically imbedded in any small portion of a Euclidean space \mathbb{R}^N , where

$$N = (n + 1)\frac{1}{2}n(3n + 11).$$

The original proof due to Nash relying on an iteration scheme has been considerably simplified. For an overview, see [?].

References

- [1] Nash, J. F., *The imbedding problem for Riemannian manifold*, Ann. of Math. 63 (1956), 20–63 (MR 17, 782)
- [2] D. Yang, *Gunther's proof of Nash's isometric embedding theorem*, <http://www.math.poly.edu/~yang/papers/gunther.pdf>online