



Math for the people, by the people.

Cartan structural equations

Canonical name	CartanStructuralEquations
Date of creation	2013-03-22 17:35:46
Last modified on	2013-03-22 17:35:46
Owner	juanman (12619)
Last modified by	juanman (12619)
Numerical id	12
Author	juanman (12619)
Entry type	Result
Classification	msc 53A45
Classification	msc 58A12
Classification	msc 58A10

To deduce the Cartan structural equations in a coordinated frame we are going to use the definition of the Christoffel symbols (connection coefficients) and where we always are going to use the Einstein sum convention:

$$\nabla_{\partial_i} \partial_j = \Gamma^s_{ij} \partial_s$$

and the curvature tensor

$$R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z$$

where X, Y, Z are any three vector fields in a riemannian manifold \mathcal{M} with the Levi-Civita connection ∇ .

First, we define through the relation $\nabla_X \partial_i = \omega^s_i(X) \partial_s$ a set of scalar function ω^s_i which are easily to see that they actually are 1-forms. We observe that $\omega^s_i(\partial_j) = \Gamma^s_{ij}$.

They satisfy skew-symmetry rule: $\omega_{si} = -\omega_{is}$, which arises from the covariant constancy of the metric tensor g_{kl} i.e.

$$\begin{aligned} 0 &= \nabla_X g_{kl} \\ &= \nabla_X \langle \partial_k, \partial_l \rangle \\ &= \langle \nabla_X \partial_k, \partial_l \rangle + \langle \partial_k, \nabla_X \partial_l \rangle \\ &= \langle \omega^s_k(X) \partial_s, \partial_l \rangle + \langle \partial_k, \omega^s_l(X) \partial_s \rangle \\ &= \omega^s_k(X) g_{sl} + \omega^s_l(X) g_{ks} \\ 0 &= \omega_{lk}(X) + \omega_{kl}(X) \end{aligned}$$

that last equation is valid for each vector field X , then $\omega_{lk} = -\omega_{kl}$.

Next we define through the relation

$$R(X, Y) \partial_i = \Omega^s_i(X, Y) \partial_s$$

the scalars $\Omega^s_i(X, Y)$ which are the so called connection 2-forms. That they are really 2-forms is an easy caligraphic exercise.

Now by the use of the Riemann curvature tensor above we see

$$\begin{aligned} R(X, Y) \partial_i &= \nabla_X \nabla_Y \partial_i - \nabla_Y \nabla_X \partial_i - \nabla_{[X, Y]} \partial_i \\ &= \nabla_X (\omega^s_i(Y) \partial_s) - \nabla_Y (\omega^s_i(X) \partial_s) - \omega^s_i[X, Y] \partial_s \\ &= X(\omega^s_i(Y)) \partial_s + \omega^s_i(Y) \nabla_X \partial_s - Y(\omega^s_i(X) \partial_s - \omega^s_i(X) \nabla_Y \partial_s - \omega^s_i[X, Y] \partial_s) \\ &= X(\omega^s_i(Y)) \partial_s + \omega^s_i(Y) \omega^t_s(X) \partial_t - Y(\omega^s_i(X) \partial_s - \omega^s_i(X) \omega^t_s(Y) \partial_t - \omega^s_i[X, Y] \partial_s) \\ &= [X(\omega^s_i(Y)) + \omega^t_i(Y) \omega^s_t(X) - Y(\omega^s_i(X)) - \omega^t_i(X) \omega^s_t(Y) - \omega^s_i[X, Y]] \partial_s \\ \Omega^s_i(X, Y) \partial_s &= [X(\omega^s_i(Y)) - Y(\omega^s_i(X)) - \omega^s_i[X, Y] + \omega^s_t(X) \omega^t_i(Y) - \omega^s_t(Y) \omega^t_i(X)] \partial_s \end{aligned}$$

In this last relation we recognize -in the first three terms- the exterior derivative of ω^s_i evaluated at (X, Y) i.e.

$$d\omega^s_i(X, Y) = X(\omega^s_i(Y)) - Y(\omega^s_i(X)) - \omega^s_i[X, Y]$$

and in the last two terms its wedge product

$$\omega^s_t \wedge \omega^t_i(X, Y) = \omega^s_t(X)\omega^t_i(Y) - \omega^s_t(Y)\omega^t_i(X)$$

all these for any two fields X, Y . Hence

$$\Omega^s_i = d\omega^s_i + \omega^s_t \wedge \omega^t_i$$

which is called the second Cartan structural equation for the coordinated frame field ∂_i .

More interesting things happen in an an-holonomic basis.