



Math for the people, by the people.

curvature of a circle

Canonical name	CurvatureOfACircle
Date of creation	2013-03-22 15:50:30
Last modified on	2013-03-22 15:50:30
Owner	cvalente (11260)
Last modified by	cvalente (11260)
Numerical id	9
Author	cvalente (11260)
Entry type	Example
Classification	msc 53A04
Related topic	circle
Related topic	curvature
Related topic	Connection
Related topic	CircleOfCurvature

Let C_r be a circle of radius r centered at the origin.

A canonical parameterization of the curve is (counterclockwise)

$$g(s) = r \left(\cos \left(\frac{s}{r} \right), \sin \left(\frac{s}{r} \right) \right)$$

for $s \in (0, 2\pi r)$ (actually this leaves out the point $(r, 0)$ but this could be treated via another parameterization taking $s \in (-\pi r, \pi r)$)

Differentiating the parameterization we get

$$\mathbf{T} = g'(s) = \left(-\sin \left(\frac{s}{r} \right), \cos \left(\frac{s}{r} \right) \right)$$

and this results in the normal

$$\mathbf{N} = J \cdot \mathbf{T} = - \left(\cos \left(\frac{s}{r} \right), \sin \left(\frac{s}{r} \right) \right) = -\frac{g(s)}{r}$$

Differentiating g a second time we can calculate the curvature

$$\mathbf{T}' = -\frac{1}{r} \left(\cos \left(\frac{s}{r} \right), \sin \left(\frac{s}{r} \right) \right) = \frac{1}{r} \mathbf{N}$$

and by definition

$$\mathbf{T}' = k\mathbf{N} \quad \therefore \quad k = \frac{1}{r}$$

and thus the curvature of a circle of radius r is $\frac{1}{r}$ provided that the positive direction on the circle is anticlockwise; otherwise it is $-\frac{1}{r}$.