

## some examples of universal bundles

Canonical name SomeExamplesOfUniversalBundles

Date of creation 2013-03-22 13:12:05 Last modified on 2013-03-22 13:12:05

Owner bwebste (988) Last modified by bwebste (988)

Numerical id 11

Author bwebste (988)
Entry type Example
Classification msc 55R15
Classification msc 55R10

Synonym universal family of spaces
Related topic CategoryOfQuantumAutomata

Defines Hilbert bundle

The universal bundle for a topological group G is usually written as  $\pi: EG \to BG$ . Any principal G-bundle for which the total space is contractible is universal; this will help us to find universal bundles without worrying about Milnor's construction of EG involving infinite joins.

- $G = \mathbb{Z}_2$ :  $E\mathbb{Z}_2 = S^{\infty}$  and  $B\mathbb{Z}_2 = \mathbb{R}P^{\infty}$ .
- $G = \mathbb{Z}_n$ :  $E\mathbb{Z}_n = S^{\infty}$  and  $B\mathbb{Z}_n = S^{\infty}/\mathbb{Z}_n$ . Here  $\mathbb{Z}_n$  acts on  $S^{\infty}$  (considered as a subset of a separable complex Hilbert space) via multiplication with an n-th root of unity.
- $G = \mathbb{Z}^n$ :  $E\mathbb{Z}^n = \mathbb{R}^n$  and  $B\mathbb{Z}^n = T^n$ .
- More generally, if G is any discrete group then one can take BG to be any Eilenberg-Mac Lane space K(G,1) and EG to be its universal cover. Indeed EG is simply connected, and it follows from the lifting theorem that  $\pi_n(EG) = 0$  for  $n \geq 0$ . This example includes the previous three and many more.
- $G = S^1$ :  $ES^1 = S^{\infty}$  and  $BS^1 = \mathbb{C}P^{\infty}$ .
- G = SU(2):  $ESU(2) = S^{\infty}$  and  $BSU(2) = \mathbb{H}P^{\infty}$ .
- G = O(n), the *n*-th orthogonal group:  $EO(n) = V(\infty, n)$ , the manifold of frames of *n* orthonormal vectors in  $\mathbb{R}^{\infty}$ , and  $BO(n) = G(\infty, n)$ , the Grassmanian of *n*-planes in  $\mathbb{R}^{\infty}$ . The projection map is taking the subspace spanned by a frame of vectors.