



Math for the people, by the people.

Brouwer fixed point theorem

Canonical name	BrouwerFixedPointTheorem
Date of creation	2013-03-22 12:44:34
Last modified on	2013-03-22 12:44:34
Owner	mathcam (2727)
Last modified by	mathcam (2727)
Numerical id	7
Author	mathcam (2727)
Entry type	Theorem
Classification	msc 55M20
Classification	msc 54H25
Classification	msc 47H10
Related topic	FixedPoint
Related topic	SchauderFixedPointTheorem
Related topic	TychonoffFixedPointTheorem
Related topic	KKMlemma
Related topic	KKMLemma

Theorem Let $\mathbf{B} = \{x \in \mathbb{R}^n : \|x\| \leq 1\}$ be the closed unit ball in \mathbb{R}^n . Any continuous function $f : \mathbf{B} \rightarrow \mathbf{B}$ has a fixed point.

Notes

Shape is not important The theorem also applies to anything homeomorphic to a closed disk, of course. In particular, we can replace \mathbf{B} in the formulation with a square or a triangle.

Compactness counts (a) The theorem is not true if we drop a point from the interior of \mathbf{B} . For example, the map $f(\vec{x}) = \frac{1}{2}\vec{x}$ has the single fixed point at 0; dropping it from the domain yields a map with no fixed points.
<http://planetmath.org/FixedPoint>

Compactness counts (b) The theorem is not true for an open disk. For instance, the map $f(\vec{x}) = \frac{1}{2}\vec{x} + (\frac{1}{2}, 0, \dots, 0)$ has its single fixed point on the boundary of \mathbf{B} .