



planetmath.org

Math for the people, by the people.

category of groupoids

Canonical name	CategoryOfGroupoids
Date of creation	2013-03-22 19:15:54
Last modified on	2013-03-22 19:15:54
Owner	bci1 (20947)
Last modified by	bci1 (20947)
Numerical id	16
Author	bci1 (20947)
Entry type	Topic
Classification	msc 55U05
Classification	msc 55U35
Classification	msc 55U40
Classification	msc 18G55
Classification	msc 18B40
Related topic	GroupoidCategory
Related topic	HomotopyCategory

1 Category of Groupoids

1.1 Properties

The category of groupoids, G_{pd} , has several important properties distinct from those of the category of groups, G_p , although it does contain the category of groups as a full subcategory. One such important property is that G_{pd} is *cartesian closed*. Thus, if J and K are two groupoids, one can form a groupoid $GPD(J, K)$ such that if G also is a groupoid then there exists a natural equivalence

$$G_{pd}(G \times J, K) \rightarrow G_{pd}(G, GPD(J, K))$$

Other important properties of G_{pd} are:

1. The category G_{pd} also has a unit interval object I , which is the groupoid with two objects $0, 1$ and exactly one arrow $0 \rightarrow 1$;
2. The groupoid I has allowed the development of a useful <http://planetmath.org/http://planetmath.org/groupoidtheory> Theory for groupoids that leads to analogies between groupoids and spaces or manifolds; effectively, groupoids may be viewed as “adding the spatial notion of a ‘place’ or location” to that of a group;
3. Groupoids extend the notion of invertible operation by comparison with that available for groups; such invertible operations also occur in the theory of inverse semigroups. Moreover, there are interesting relations between inverse semigroups and ordered groupoids. Such concepts are thus applicable to sequential machines and automata whose state spaces are semigroups. Interestingly, the category of finite automata, just like G_{pd} is also *cartesian closed*;
4. The category G_{pd} has a variety of types of morphisms, such as: quotient morphisms, retractions, covering morphisms, fibrations, universal morphisms, (in contrast to only the epimorphisms and monomorphisms of group theory);
5. A monoid object, $END(J) = GPD(J, J)$, also exists in the category of groupoids, that contains a maximal subgroup object denoted here as $AUT(J)$. Regarded as a group object in the category groupoids,

$AUT(J)$ is equivalent to a crossed module C_M , which in the case when J is a group is the traditional crossed module $J \rightarrow Aut(J)$, defined by the inner automorphisms.

References

- [1] May, J.P. 1999, *A Concise Course in Algebraic Topology.*, The University of Chicago Press: Chicago
- [2] R. Brown and G. Janelidze.(2004). Galois theory and a new homotopy double groupoid of a map of spaces.(2004). *Applied Categorical Structures*,**12**: 63-80. Pdf file in arxiv: math.AT/0208211
- [3] P. J. Higgins. 1971. *Categories and Groupoids.*, Originally published by: Van Nostrand Reinhold, 1971 Republished in: *Reprints in Theory and Applications of Categories*, No. 7 (2005) pp 1-195: <http://www.tac.mta.ca/tac/reprints/articles/7/tr7.pdf>