



**planetmath.org**

Math for the people, by the people.

## (path) connectness as a homotopy invariant

Canonical name	pathConnectnessAsAHomotopyInvariant
Date of creation	2013-03-22 18:02:15
Last modified on	2013-03-22 18:02:15
Owner	joking (16130)
Last modified by	joking (16130)
Numerical id	8
Author	joking (16130)
Entry type	Theorem
Classification	msc 55P10
Related topic	Homotopy
Related topic	homotopyequivalence
Related topic	path
Related topic	connectedspace

**Theorem.** Let  $X$  and  $Y$  be arbitrary topological spaces with  $Y$  (path) connected. If there are maps  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$  such that  $g \circ f : X \rightarrow X$  is homotopic to the identity map, then  $X$  is (path) connected.

**Proof:** Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$  be maps satisfying theorem's assumption. Furthermore let  $X = \bigcup X_i$  be a decomposition of  $X$  into (path) connected components. Since  $Y$  is (path) connected, then  $g(Y) \subseteq X_i$  for some  $i$ . Thus  $(g \circ f)(X) \subseteq X_i$ . Now let  $H : I \times X \rightarrow X$  be the homotopy from  $g \circ f$  to the identity map. Let  $\alpha_x : I \rightarrow X$  be a path defined by the formula:  $\alpha_x(t) = H(t, x)$ . Since for all  $x \in X$  we have  $\alpha_x(0) \in X_i$  and  $I$  is path connected, then  $\alpha_x(I) \subseteq X_i$ . Therefore  $H(I \times X) \subseteq X_i$ , but  $H(\{1\} \times X) = X$  which implies that  $X_i = X$ , so  $X$  is (path) connected.  $\square$

Straightforward application of this theorem is following:

**Corollary.** Let  $X$  and  $Y$  be homotopy equivalent spaces. Then  $X$  is (path) connected if and only if  $Y$  is (path) connected.