



planetmath.org

Math for the people, by the people.

relative homology groups

Canonical name	RelativeHomologyGroups
Date of creation	2013-03-22 13:14:47
Last modified on	2013-03-22 13:14:47
Owner	bwebste (988)
Last modified by	bwebste (988)
Numerical id	5
Author	bwebste (988)
Entry type	Definition
Classification	msc 55N10

If X is a topological space, and A a subspace, then the inclusion map $A \hookrightarrow X$ makes $C_n(A)$ into a subgroup of $C_n(X)$. Since the boundary map on $C_*(X)$ restricts to the boundary map on $C_*(A)$, we can take the quotient complex $C_*(X, A)$,

$$\xleftarrow{\partial} C_n(X)/C_n(A) \xleftarrow{\partial} C_{n+1}(X)/C_{n+1}(A) \xleftarrow{\partial}$$

The homology groups of this complex $H_n(X, A)$, are called the *relative homology groups* of the pair (X, A) . Under relatively mild hypotheses, $H_n(X, A) = H_n(X/A)$ where X/A is the set of equivalence classes of the relation $x \sim y$ if $x = y$ or if $x, y \in A$, given the quotient topology (this is essentially X , with A reduced to a single point). Relative homology groups are important for a number of reasons, principally for computational ones, since they fit into long exact sequences, which are powerful computational tools in homology.