



Math for the people, by the people.

Sard's theorem

Canonical name	SardsTheorem
Date of creation	2013-03-22 13:04:09
Last modified on	2013-03-22 13:04:09
Owner	mathcam (2727)
Last modified by	mathcam (2727)
Numerical id	9
Author	mathcam (2727)
Entry type	Theorem
Classification	msc 57R35
Related topic	Residual
Related topic	BaireCategoryTheorem
Defines	critical point
Defines	critical value
Defines	regular value

Let $\phi : X^n \rightarrow Y^m$ be a smooth map on smooth manifolds. A *critical point* of ϕ is a point $p \in X$ such that the differential $\phi_* : T_p X \rightarrow T_{\phi(p)} Y$ considered as a linear transformation of real vector spaces has <http://planetmath.org/RankLinearMapping> $< m$. A *critical value* of ϕ is the image of a critical point. A *regular value* of ϕ is a point $q \in Y$ which is not the image of any critical point. In particular, q is a regular value of ϕ if $q \in Y \setminus \phi(X)$.

Following Spivak [?], we say a subset V of Y^m *has measure zero* if there is a sequence of coordinate charts (x_i, U_i) whose union contains V and such that $x_i(U_i \cap V)$ has measure 0 (in the usual sense) in \mathbb{R}^m for all i . With that definition, we can now state:

Sard's Theorem. Let $\phi : X \rightarrow Y$ be a smooth map on smooth manifolds. Then the set of critical values of ϕ has measure zero.

References

- [Spivak] Spivak, Michael. *A Comprehensive Introduction to Differential Geometry*. Volume I, Third Edition. Publish of Perish, Inc. Houston, Texas. 1999.