



planetmath.org

Math for the people, by the people.

Brouwer degree

Canonical name	BrouwerDegree
Date of creation	2013-03-22 14:52:37
Last modified on	2013-03-22 14:52:37
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	7
Author	jirka (4157)
Entry type	Definition
Classification	msc 57R35
Synonym	degree
Related topic	DegreeMod2OfAMapping

Suppose that M and N are two oriented differentiable manifolds of dimension n (without boundary) with M compact and N connected and suppose that $f: M \rightarrow N$ is a differentiable mapping. Let $Df(x)$ denote the differential mapping at the point $x \in M$, that is the linear mapping $Df(x): T_x(M) \rightarrow T_{f(x)}(N)$. Let $\text{sign } Df(x)$ denote the sign of the determinant of $Df(x)$. That is the sign is positive if f preserves orientation and negative if f reverses orientation.

Definition. Let $y \in N$ be a regular value, then we define the *Brower degree* (or just degree) of f by

$$\deg f := \sum_{x \in f^{-1}(y)} \text{sign } Df(x).$$

It can be shown that the degree does not depend on the regular value y that we pick so that $\deg f$ is well defined.

Note that this degree coincides with the <http://planetmath.org/Degree5degree> as defined for maps of spheres.

References

- [1] John W. Milnor. . The University Press of Virginia, Charlottesville, Virginia, 1969.