



planetmath.org

Math for the people, by the people.

loop theorem

Canonical name	LoopTheorem
Date of creation	2013-03-22 15:49:13
Last modified on	2013-03-22 15:49:13
Owner	juanman (12619)
Last modified by	juanman (12619)
Numerical id	11
Author	juanman (12619)
Entry type	Theorem
Classification	msc 57M35
Related topic	3Manifolds

In the topology of 3-manifolds, **the loop theorem** is generalization of an ansatz discovered by Max Dehn (namely, Dehn's lemma), who saw that *if a continuous map from a 2-disk to a 3-manifold whose restriction to the boundary's disk has no singularities, then there exists another embedding whose restriction to the boundary's disk is equal to the boundary's restriction original map.*

The following statement called the loop theorem is a version from J. Stallings, but written in W. Jaco's book.

*Let M be a three-manifold and let S be a connected surface in ∂M . Let $N \subset \pi_1(M)$ be a normal subgroup. Let $f: D^2 \rightarrow M$ be a **continuous map** such that $f(\partial D^2) \subset S$ and $[f|\partial D^2] \notin N$.*

*Then there exists an **embedding** $g: D^2 \rightarrow M$ such that $g(\partial D^2) \subset S$ and $[g|\partial D^2] \notin N$,*

The proof is a clever construction due to C. Papakyriakopoulos about a sequence (a tower) of covering spaces. Maybe the best detailed presentation is due to A. Hatcher. But in general, accordingly to Jaco's opinion, "*... for anyone unfamiliar with the techniques of 3-manifold-topology and are here to gain a working knowledge for the study of problems in this ..., there is no better to start.*"

References

W. Jaco, *Lectures on 3-manifolds topology*, A.M.S. regional conference series in Math 43.

J. Hempel, *3-manifolds*, Princeton University Press 1976.

A. Hatcher, *Notes on 3-manifolds*, available on-line.