



Math for the people, by the people.

Finsler geometry

Canonical name	FinslerGeometry
Date of creation	2013-03-22 15:03:37
Last modified on	2013-03-22 15:03:37
Owner	paolini (1187)
Last modified by	paolini (1187)
Numerical id	8
Author	paolini (1187)
Entry type	Definition
Classification	msc 58B20
Classification	msc 53B40
Classification	msc 53C60
Related topic	WulffTheorem

Let \mathcal{M} be an n -dimensional differential manifold and let $\phi: T\mathcal{M} \rightarrow \mathbb{R}$ be a function $\phi(x, \xi)$ defined for $x \in \mathcal{M}$ and $\xi \in T_x\mathcal{M}$ such that $\phi(x, \cdot)$ is a possibly non symmetric norm on $T_x\mathcal{M}$. The couple (\mathcal{M}, ϕ) is called a Finsler space.

Let us define the ϕ -length of curves in \mathcal{M} . If $\gamma: [a, b] \rightarrow \mathcal{M}$ is a differentiable curve we define

$$\ell_\phi(\gamma) := \int_a^b \phi(\gamma'(t)) dt.$$

So a natural geodesic distance can be defined on \mathcal{M} which turns the Finsler space into a quasi-metric space (if \mathcal{M} is connected):

$$d_\phi(x, y) := \inf\{\ell_\phi(\gamma) : \gamma \text{ is a differentiable curve } \gamma: [a, b] \rightarrow \mathcal{M} \text{ such that } \gamma(a) = x \text{ and } \gamma(b) = y\}.$$

Notice that every Riemann manifold (\mathcal{M}, g) is also a Finsler space, the norm $\phi(x, \cdot)$ being the norm induced by the scalar product $g(x)$.

A finite dimensional Banach space is another simple example of Finsler space, where $\phi(x, \xi) := \|\xi\|$. Wulff Theorem is one of the most important theorems in this ambient space.