



Math for the people, by the people.

normal bundle

Canonical name	NormalBundle
Date of creation	2013-03-22 13:59:06
Last modified on	2013-03-22 13:59:06
Owner	bwebste (988)
Last modified by	bwebste (988)
Numerical id	5
Author	bwebste (988)
Entry type	Definition
Classification	msc 58A32

Let  $X$  be an immersed submanifold of  $M$ , with immersion  $i : X \rightarrow M$ . Then we can restrict the tangent bundle of  $M$  to  $N$  or more properly, take the pullback  $i^*TM$ . This, as a vector bundle over  $X$  should contain a lot of information about the embedding of  $X$  into  $M$ . But there is a natural injection  $TX \rightarrow i^*TM$ , and the subbundle which is the image of this only has information on the intrinsic properties of  $X$ , and thus is useless in obtaining information about the embedding of  $X$  into  $M$ . Instead, to get information on this, we take the quotient  $i^*TM/TX = NX$ , the normal bundle of  $X$ . The normal bundle is very strongly dependent on the immersion  $i$ . If  $E$  is any vector bundle on  $X$ , then  $E$  is the normal bundle for the embedding of  $X$  into  $E$  as the zero section.

The normal bundle determines the local geometry of the embedding of  $X$  into  $M$  in the following sense: In  $M$ , there exists an open neighborhood  $U \supset X$  which is diffeomorphic to  $NX$  by a diffeomorphism taking  $X$  to the zero section.