



planetmath.org

Math for the people, by the people.

differential form

Canonical name	DifferentialForm
Date of creation	2013-03-22 12:44:46
Last modified on	2013-03-22 12:44:46
Owner	rmilson (146)
Last modified by	rmilson (146)
Numerical id	28
Author	rmilson (146)
Entry type	Definition
Classification	msc 58A10
Defines	exterior derivative
Defines	1-form
Defines	exterior product
Defines	wedge product
Defines	interior product
Defines	tensorial

# 1 Notation and Preliminaries.

Let  $M$  be an  $n$ -dimensional differential manifold. Let  $TM$  denote the manifold's tangent bundle,  $C^\infty(M)$  the algebra of smooth functions, and  $V(M)$  the Lie algebra of smooth vector fields. The directional derivative makes  $C^\infty(M)$  into a  $V(M)$  module. Using local coordinates, the directional derivative operation can be expressed as

$$v(f) = v^i \partial_i f, \quad v \in V(M), \quad f \in C^\infty(M).$$

## 2 Definitions.

**Differential forms.** Let  $A$  be a  $C^\infty(M)$  module. An  $\mathbb{R}$ -linear mapping  $\alpha : V(M) \rightarrow A$  is said to be *tensorial* if it is a  $C^\infty(M)$ -homomorphism, in other words, if it satisfies

$$\alpha(fv) = f\alpha(v)$$

for all vector fields  $v \in V(M)$  and functions  $f \in C^\infty(M)$ . More generally, a multilinear map  $\alpha : V(M) \times \cdots \times V(M) \rightarrow A$  is called tensorial if it satisfies

$$\alpha(fu, \dots, v) = \cdots = \alpha(u, \dots, fv) = f\alpha(u, \dots, v)$$

for all vector fields  $u, \dots, v$  and all functions  $f \in C^\infty(M)$ .

We now define a differential 1-form to be a tensorial linear mapping from  $V(M)$  to  $C^\infty(M)$ . More generally, for  $k = 0, 1, 2, \dots$ , we define a differential  $k$ -form to be a tensorial multilinear, antisymmetric, mapping from  $V(M) \times \cdots \times V(M)$  ( $k$  times) to  $C^\infty(M)$ . Using slightly fancier language, the above amounts to saying that a 1-form is a section of the cotangent bundle  $T^*M = \text{Hom}(TM, \mathbb{R})$ , while a differential  $k$ -form is a section of  $\text{Hom}(\Lambda^k TM, \mathbb{R})$ .

Henceforth, we let  $\Omega^k(M)$  denote the  $C^\infty(M)$ -module of differential  $k$ -forms. In particular, a differential 0-form is the same thing as a function. Since the tangent spaces of  $M$  are  $n$ -dimensional vector spaces, we also have  $\Omega^k(M) = 0$  for  $k > n$ . We let

$$\Omega(M) = \bigoplus_{k=0}^n \Omega^k(M)$$

denote the vector space of all differential forms. There is a natural operator, called the exterior product, that endows  $\Omega(M)$  with the structure of a graded algebra. We describe this operation below.

**Exterior and Interior Product.** Let  $v \in V(M)$  be a vector field and  $\alpha \in \Omega^k(M)$  a differential form. We define  $\iota_v(\omega)$ , the interior product of  $v$  and  $\alpha$ , to be the differential  $k-1$  form given by

$$\iota_v(\alpha)(u_1, \dots, u_{k-1}) = \alpha(v, u_1, \dots, u_{k-1}), \quad v_1, \dots, v_{k-1} \in V(M).$$

The interior product of a vector field with a 0-form is defined to be zero.

Let  $\alpha \in \Omega^k(M)$  and  $\beta \in \Omega^\ell(M)$  be differential forms. We define the exterior, or wedge product  $\alpha \wedge \beta \in \Omega^{k+\ell}(M)$  to be the unique differential form such that

$$\iota_v(\alpha \wedge \beta) = \iota_v(\alpha) \wedge \beta + (-1)^k \alpha \wedge \iota_v(\beta)$$

for all vector fields  $v \in V(M)$ . Equivalently, we could have defined

$$(\alpha \wedge \beta)(v_1, \dots, v_{k+\ell}) = \sum_{\pi} \text{sgn}(\pi) \alpha(v_{\pi_1}, \dots, v_{\pi_k}) \beta(v_{\pi_{k+1}}, \dots, v_{\pi_{k+\ell}}),$$

where the sum is taken over all permutations  $\pi$  of  $\{1, 2, \dots, k+\ell\}$  such that  $\pi_1 < \pi_2 < \dots < \pi_k$  and  $\pi_{k+1} < \dots < \pi_{k+\ell}$ , and where  $\text{sgn} \pi = \pm 1$  according to whether  $\pi$  is an even or odd permutation.

**Exterior derivative.** The exterior derivative is a first-order differential operator  $d : \Omega^*(M) \rightarrow \Omega^*(M)$ , that can be defined as the unique linear mapping satisfying

$$\begin{aligned} d(d\alpha) &= 0, & \alpha &\in \Omega^k(M); \\ \iota_V(df) &= v(f), & v &\in V(M), f \in C^\infty(M); \\ d(\alpha \wedge \beta) &= d(\alpha) \wedge \beta + (-1)^k \alpha \wedge d(\beta), & \alpha &\in \Omega^k(M), \beta \in \Omega^\ell(M). \end{aligned}$$

### 3 Local coordinates.

Let  $(x^1, \dots, x^n)$  be a system of local coordinates on  $M$ , and let  $\partial_1, \dots, \partial_n$  denote the corresponding frame of coordinate vector fields. In other words,

$$\partial_i(x^j) = \delta_i^j,$$

where the right hand side is the usual Kronecker delta symbol. By the definition of the exterior derivative,

$$\iota_{\partial_i}(dx^j) = \delta_i^j;$$

In other words, the 1-forms  $dx^1, \dots, dx^n$  form the dual coframe.

Locally, the  $\partial_i$  freely generate  $V(M)$ , meaning that every vector field  $v \in V(M)$  has the form

$$v = v^i \partial_i,$$

where the coordinate components  $v^i$  are uniquely determined as

$$v^i = v(x^i).$$

Similarly, locally the  $dx^i$  freely generate  $\Omega^1(M)$ . This means that every one-form  $\alpha \in \Omega^1(M)$  takes the form

$$\alpha = \alpha_i dx^i,$$

where

$$\alpha_i = \iota_{\partial_i}(\alpha).$$

More generally, locally  $\Omega^k(M)$  is a freely generated by the differential  $k$ -forms

$$dx^{i_1} \wedge \dots \wedge dx^{i_k}, \quad 1 \leq i_1 < i_2 < \dots < i_k \leq n.$$

Thus, a differential form  $\alpha \in \Omega^k(M)$  is given by

$$\begin{aligned} \alpha &= \sum_{i_1 < \dots < i_k} \alpha_{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k}, \\ &= \frac{1}{k!} \alpha_{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k}, \end{aligned} \tag{1}$$

where

$$\alpha_{i_1 \dots i_k} = \alpha(\partial_{i_1}, \dots, \partial_{i_k}).$$

Consequently, for vector fields  $u, v, \dots, w \in V(M)$ , we have

$$\alpha(u, v, \dots, w) = \alpha_{i_1 i_2 \dots i_k} u^{i_1} v^{i_2} \dots w^{i_k}.$$

In terms of local coordinates and the skew-symmetrization index notation, the interior and exterior product, and the exterior derivative take the following expressions:

$$(\iota_v(\alpha))_{i_1 \dots i_k} = v^j \alpha_{j i_1 \dots i_k}, \quad v \in V(M), \alpha \in \Omega^{k+1}(M); \tag{2}$$

$$(\alpha \wedge \beta)_{i_1 \dots i_{k+\ell}} = \binom{k+\ell}{k} \alpha_{[i_1 \dots i_k} \beta_{i_{k+1} \dots i_{k+\ell}]}, \quad \alpha \in \Omega^k(M), \beta \in \Omega^\ell(M); \tag{3}$$

$$(d\alpha)_{i_0 i_1 \dots i_k} = (k+1) \partial_{[i_0} \alpha_{i_1 \dots i_k]}, \quad \alpha \in \Omega^k(M). \tag{4}$$

Note that some authors prefer a different definition of the components of a differential. According to this alternate convention, a factor of  $k!$  placed before the summation sign in (??), and the leading factors are removed from (??) and (??).