

planetmath.org

Math for the people, by the people.

volume element

Canonical name VolumeElement
Date of creation 2013-03-22 17:40:58
Last modified on 2013-03-22 17:40:58

Owner jirka (4157) Last modified by jirka (4157)

Numerical id 5

Author jirka (4157) Entry type Definition Classification msc 58A10msc 53-00Classification Synonym volume form Synonym volume measure Defines area element Defines area form Defines area measure

Defines Euclidean volume element
Defines Euclidean volume form
Defines euclidean volume measure

Defines surface area measure
Defines surface area element
Defines surface area form

If M is an n dimensional manifold, then a http://planetmath.org/DifferentialFormsdifferent n form that is never zero is called a *volume element* or a *volume form*. Usually one volume form is associated with the manifold. The volume element is sometimes denoted by dV, ω or vol_n . If the manifold is a Riemannian manifold with g, then the natural volume form is defined in local coordinates $x^1 \dots x^n$ by

 $dV := \sqrt{|g|} dx^1 \wedge \ldots \wedge dx^n.$

It is not hard to show that a manifold has a volume form if and only if it is orientable.

If the manifold is \mathbb{R}^n , then the usual volume element $dV = dx^1 \wedge dx^2 \wedge \ldots \wedge dx^n$ is called the *Euclidean volume element* or *Euclidean volume form*. In this context, \mathbb{C}^n is usually treated as \mathbb{R}^{2n} unless stated otherwise.

When n=2, then the form is frequently called the *area element* or *area* form and frequently denoted by dA. Furthermore, when the manifold is a submanifold of \mathbb{R}^3 , then many authors will refer to a surface area element or surface area form.

When the context is measure theoretic, this form is sometimes called a volume measure, area measure, etc...

References

- [1] Michael Spivak., W.A. Benjamin, Inc., 1965.
- [2] William M. Boothby. , Academic Press, San Diego, California, 2003.