



Math for the people, by the people.

## Cochran's theorem

Canonical name	CochransTheorem
Date of creation	2013-03-22 14:33:01
Last modified on	2013-03-22 14:33:01
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	8
Author	CWoo (3771)
Entry type	Theorem
Classification	msc 62J10
Classification	msc 62H10
Classification	msc 62E10
Defines	Fisher's theorem

Let  $\mathbf{X}$  be multivariate normally distributed as  $\mathbf{N}_p(\mathbf{0}, \mathbf{I})$  such that

$$\mathbf{X}^T \mathbf{X} = \sum_{i=1}^k Q_i,$$

where each

1.  $Q_i$  is a quadratic form
2.  $Q_i = \mathbf{X}^T \mathbf{B}_i \mathbf{X}$ , where  $\mathbf{B}_i$  is a  $p$  by  $p$  square matrix
3.  $\mathbf{B}_i$  is positive semidefinite
4.  $\text{rank}(\mathbf{B}_i) = r_i$

Then any two of the following imply the third:

1.  $\sum_{i=1}^k r_i = p$
2. each  $Q_i$  has a <http://planetmath.org/ChiSquaredRandomVariablechi> square distribution with  $r_i$  of freedom,  $\chi^2(r_i)$
3.  $Q_i$ 's are mutually independent

As an example, suppose  $X_1^2 \sim \chi^2(m_1)$  and  $X_2^2 \sim \chi^2(m_2)$ . Furthermore, assume  $X_1^2 \geq X_2^2$  and  $m_1 > m_2$ , then

$$X_1^2 - X_2^2 \sim \chi^2(m_1 - m_2).$$

This corollary is known as *Fisher's theorem*.