

planetmath.org

Math for the people, by the people.

Frobenius matrix norm

Canonical name FrobeniusMatrixNorm
Date of creation 2013-03-22 11:43:25
Last modified on 2013-03-22 11:43:25
Owner mathcam (2727)
Last modified by mathcam (2727)

Numerical id 25

Author mathcam (2727)

Entry type Definition
Classification msc 65F35
Classification msc 15A60
Classification msc 18-00

Synonym Euclidean matrix norm

Synonym matrix F-norm

Synonym Hilbert-Schmidt norm

Related topic MatrixNorm
Related topic MatrixPnorm
Related topic VectorNorm
Related topic VectorPnorm
Related topic ShursInequality

Related topic trace
Related topic transpose
Related topic Transpose

Related topic MatrixLogarithm Related topic FrobeniusProduct Let R be a ring with a valuation $|\cdot|$ and let M(R) denote the set of matrices over R. The Frobenius norm function or Euclidean matrix norm is the norm function $||\cdot||_F: M(R) \to \mathbb{R}$ given by

$$||A||_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^2},$$

where m and n respectively denote the number of rows and columns of A. Note A need not be square for this definition. A more concise (though) definition, in the case that $R = \mathbb{R}$ or \mathbb{C} , is

$$||A||_F = \sqrt{\operatorname{trace}(A^*A)},$$

where A^* denotes the conjugate transpose of A.

Some:

• Denote the columns of A by A_i . A nice property of the norm is that

$$||A||_F^2 = ||A_1||_2^2 + ||A_2||_2^2 + \dots + ||A_n||_2^2.$$

- Let A be a square matrix and let U be a unitary matrix of same size as A. Then $||A||_F = ||U^*AU||_F$ where U^* is the conjugate transpose of U.
- If AB is defined, then $||AB||_F \le ||A||_F ||B||_F$.