



planetmath.org

Math for the people, by the people.

QR decomposition

Canonical name	QRDecomposition
Date of creation	2013-03-22 12:06:04
Last modified on	2013-03-22 12:06:04
Owner	akrowne (2)
Last modified by	akrowne (2)
Numerical id	8
Author	akrowne (2)
Entry type	Definition
Classification	msc 65F25
Classification	msc 15A23
Synonym	QR factorization
Synonym	QR-decomposition
Synonym	QR-factorization
Related topic	LUDecomposition
Related topic	GramSchmidtOrthogonalization

# 1 QR Decomposition

Orthogonal matrix triangularization (QR decomposition) reduces a real  $m \times n$  matrix  $A$  with  $m \geq n$  and full rank to a much simpler form. It guarantees numerical stability by minimizing errors caused by machine roundoffs. A suitably chosen orthogonal matrix  $Q$  will triangularize the given matrix:

$$A = Q \begin{bmatrix} R \\ 0 \end{bmatrix}$$

with the  $n \times n$  right triangular matrix  $R$ . One only has then to solve the triangular system  $Rx = Pb$ , where  $P$  consists of the first  $n$  rows of  $Q$ .

The least squares problem  $Ax \approx b$  is easy to solve with  $A = QR$  and  $Q$  an orthogonal matrix (here and henceforth  $R$  is the entire  $m \times n$  augmented matrix from above). The solution

$$x = (A^T A)^{-1} A^T b$$

becomes

$$x = (R^T Q^T Q R)^{-1} R^T Q^T b = (R^T R)^{-1} R^T Q^T b = R^{-1} Q^T b$$

This is a matrix-vector multiplication  $Q^T b$ , followed by the solution of the triangular system  $Rx = Q^T b$  by back-substitution. The QR factorization saves us the formation of  $A^T A$  and the solution of the normal equations.

Many different methods exist for the QR decomposition, e.g. the Householder transformation, the Givens rotation, or the Gram-Schmidt decomposition.

## References

- [1] The Data Analysis Briefbook. <http://rkb.home.cern.ch/rkb/titleA.html><http://rkb.home.cern.ch/rkb/titleA.html>